

---

# SILVA FENNICA

Monograph 4 · 2005

---

Dimitris Zianis, Petteri Muukkonen, Raisa Mäkipää  
and Maurizio Mencuccini

## Biomass and Stem Volume Equations for Tree Species in Europe

---

The Finnish Society of Forest Science  
The Finnish Forest Research Institute

## Abstract

---

**Zianis, D., Muukkonen, P., Mäkipää, R. & Mencuccini, M.** 2005. Biomass and stem volume equations for tree species in Europe. *Silva Fennica Monographs* 4. 63 p.

A review of stem volume and biomass equations for tree species growing in Europe is presented. The mathematical forms of the empirical models, the associated statistical parameters and information about the size of the trees and the country of origin were collated from scientific articles and from technical reports. The total number of the compiled equations for biomass estimation was 607 and for stem volume prediction it was 230. The analysis indicated that most of the biomass equations were developed for aboveground tree components. A relatively small number of equations were developed for southern Europe. Most of the biomass equations were based on a few sampled sites with a very limited number of sampled trees. The volume equations were, in general, based on more representative data covering larger geographical regions. The volume equations were available for major tree species in Europe. The collected information provides a basic tool for estimation of carbon stocks and nutrient balance of forest ecosystems across Europe as well as for validation of theoretical models of biomass allocation.

**Keywords** aboveground biomass, allometry, belowground biomass, biomass function, dbh, tree diameter, tree height

**Correspondence** Mäkipää: Finnish Forest Research Institute, Unioninkatu 40A, FIN-00170, Helsinki, Finland; Zianis & Mencuccini: Department of Forestry, TEI Kavalas, Drama 66050, Greece

**E-mail** raisa.makipaa@metla.fi

**Received** 23 March 2004 **Revised** 6 September 2005 **Accepted** 7 September 2005

---

ISBN 951-40-xxx (print), ISBN 951-40-xxx (pdf)  
ISSN 1457-7356

Printed by Tammer-Paino Oy, Tampere, Finland

# Contents

---

1 INTRODUCTION.....	5
2 MATERIAL AND METHODS .....	6
3 RESULTS.....	7
3.1 Biomass Equations.....	7
3.2 Stem Volume Equations .....	11
4 DISCUSSION .....	14
REFERENCES .....	16
Appendix A. Biomass equations for different biomass components by tree species .....	18
Appendix B. General descriptions of volume equations.....	44
Appendix C. Volume equations for different tree species.....	52

## Acknowledgments

---

The authors wish to express their thanks to the participants of the European Co-operation in the field of Scientific and Technical Research (COST) E21 Action ‘Contribution of forests and forestry to mitigate greenhouse effects’ and other scientists for submitting equations for the presented database. The study was carried out with the financial support of the Finnish Ministry of Agriculture and Forestry, and the EU-funded research consortium ‘Multi-source inventory methods for quantifying carbon stocks and stock changes in European forests’ (CarboInvent EKV2-CT-2002-00157). In addition, Dr. Dimitris Zianis was partially supported by the I.K.Y (Scholarship State Foundation of Greece), and prof. **M. Mencuccini** by the EU-funded Carbo-Age project (EVK2-1999-CT-00045).

# 1 Introduction

---

The estimation of stem volume and tree biomass is needed for both sustainable planning of forest resources and for studies on the energy and nutrients flows in ecosystems. Planners at the strategic and operational levels have strongly emphasised the need for accurate estimates of stem volume, while Hall (1997) reviewed the potential role of biomass as an energy source in the 21st century. In addition, the United Nations Framework Convention on Climate Change and in particular the Kyoto Protocol recognise the importance of forest carbon sink and the need to monitor, preserve and enhance terrestrial carbon stocks, since changes in the forest carbon stock influence the atmospheric CO<sub>2</sub> concentration. Terrestrial biotic carbon stocks and stock changes are difficult to assess (IPCC 2003) and most current estimates are subject to considerable uncertainty (Löwe et al. 2000, Clark et al. 2001, Jenkins et al. 2003). The reliability of the current estimates of the forest carbon stock and the understanding of ecosystem carbon dynamics can be improved by applying existing knowledge on the allometry of trees that is available in the form of biomass and volume equations (Jenkins et al. 2003, Zianis and Mencuccini 2003, Lehtonen et al. 2004). The biomass equations can be applied directly to tree level inventory data (the measured dimensions of trees; diameter, height), or biomass expansion factors (BEFs) applicable to stand level inventory data can be developed and tested with the help of representative volume and biomass equations (Lehtonen et al. 2004).

Recently, remote sensing data have been used to assess standing volume and forest biomass (Montes et al. 2000, Drake et al. 2002). However, the estimation of biomass depends on ground truth data with measured dimensions of trees, and the empirical biomass equations are therefore needed to predict biomass as a function of recorded variables.

The wealth of allometric equations that relate stem volume as well as the biomass of several tree components to diameter at breast height and/or to tree height has never been summarised for European tree species, although this has been for American (Tritton and Hornbeck 1982, Ter-Mikaelian and Korzukhin 1997, Jenkins et al. 2004) and Australian trees (Eamus et al. 2000, Keith et al. 2000). Since the development of stem volume and biomass equations is laborious and time consuming process – especially the destructive harvesting of large trees – existing equations need to be compiled and evaluated to facilitate identification of the gaps in the coverage of the equations. The compiled equations can also be used to test and compare existing equations with new ones as well as to validate process-based models.

The aim of this study was to develop a database on tree-level stem volume and biomass equations for various tree species growing in Europe. Equations for both whole tree biomass and the biomass of different components were considered. The compiled database is a guide to the original publications of these equations. In ecological studies on forest carbon and nutrient cycling, forest and greenhouse gas inventories as well as in the validation of process-based models, this database facilitates effective exploitation of existing information on the allometry of trees.

## 2 Material and Methods

---

The development of the presented compilation of equations was based on published equations for different tree species growing on the European continent. We restricted the compilation to the relationships published on the European continent since similar kinds of information have already been presented for different biomes (Zianis and Mencuccini 2004), for North American tree species (Ter-Mikaelian and Korzukhin 1997, Jenkins et al. 2004), and for Australian ecosystems (see reports by Eamus et al. 2000, Keith et al. 2000, Snowdon et al. 2000). In order to compile the available information we conducted a literature survey on forestry and forest-related journals. However, part of the equations, particularly for stem volume relationships, have been published in the technical reports of research institutes or research programmes across Europe. In many cases, the original papers had not been written in the English language. To obtain these equations, researchers throughout Europe were asked to provide any allometric equation published in their country and readily available to them.

For all the empirical relationships included in the database, the explanatory variables were always the diameter at breast height ( $D$ ), the tree height ( $H$ ) or a combination of the two. For latest decades, standardized reference point for breast height and height measurements has been ground level and, in the European countries, the stem diameter at breast height have been measured at 1.3 above ground (Bruce and Schumacher 1950, Köhl et al. 1997). These two variables ( $D$  and  $H$ ) are the most commonly used independent variables, but equations with several other independent variables (e.g. site fertility, elevation, soil type) have been also widely developed. Those equations were not, however, included in this database, since selection of variables it is highly dependent on local conditions and intended local use of equations. Some empirical relationships reported in the original articles were excluded

from this review and database since the equations with reported values of the parameters generated estimates that were not realistic (e.g. negative values, or shape of equation indicate impossible allometry of trees). In addition, equations with notably low  $r^2$ -values were excluded. In the original publications, there might occur several other equations besides the one compiled in the present study. No selection criteria were applied with regard to the species, age, size, site conditions, or sampling method. The compiled biomass equations were presented according to different tree components (Table 1).

The measurement units for the regressed and the explanatory variables, the number of the sampled trees ( $n$ ), the coefficient of determination ( $r^2$ ), and the range of diameter and height were also included in this review whenever this information was available in the original article. Additionally, the basal area of the stand and the stand density from which the sampled trees originated, the location (longitude and latitude) of the sampled trees as well as the standard error of the parameters of the regressions, the type and corresponding value of the statistical error, and the correction factor (Sprugel 1983) were also collected for the compiled equations. However, information on these parameters is not shown in the Appendix of the present study since it was reported only in a very limited number of original articles.

## 3 Results

---

### 3.1 Biomass Equations

We found biomass equations for various aboveground and belowground components (Table 1), but most of the biomass equations were for aboveground parts, particularly for branches and foliage (Table 2). Very few equations were available for the biomass of dead branches, coarse, small and fine roots, and only four to estimate the biomass of cones (Table 2). The total number of the compiled biomass equations for different tree components was 607 (Appendix A).

The compiled biomass equations refer to 39 different tree species growing in Europe (Table 3). The vast majority of the compiled empirical equations developed for different tree components was reported for northern and central European countries (Table 3). Totally 82 equations referred to data recorded in southern European countries, particularly Greece, Italy, Portugal and Spain.

**Table 1.** Abbreviations for tree biomass components.

AB	Total aboveground biomass
ABW	Total aboveground woody biomass
BR	Branch biomass
CO	Biomass of cones
CR	Crown biomass (BR+FL)
DB	Biomass of dead branches
FL	Total foliage biomass
FL( <i>i</i> )	Biomass of <i>i</i> -year-old needles
RC	Biomass of coarse roots <sup>a</sup>
RF	Biomass of fine roots <sup>a</sup>
RS	Biomass of small roots <sup>a</sup>
RT	Biomass of roots (RC+RF+RS)
SB	Biomass of stem bark
SR	Biomass of the stump-root system <sup>a</sup>
ST	Total stem biomass (SW+SB)
SU	Stump biomass <sup>a</sup>
SW	Stem wood biomass
TB	Total tree biomass (AB+RT)
TW	Total woody biomass

<sup>a</sup>Defined differently in each study

For the some biomass equations of *Abies balsamea* (L.) Mill., *Fagus crenata* Bl., *Picea rubens* Sarg., *Pinus banksiana* Lamb., *Pinus contorta* Doug. ex Loud., and *Pinus taeda* L., the location of the sampled trees was not reported. Only one equation was available for each of the following components: branch biomass within the crown, the biomass of epicormic branches, stem biomass within the crown, woody biomass in the crown, foliage biomass in crown, foliage biomass of epicormic branches (reported by Zianis and Mencuccini 2003). Thus, they were not included in the database.

The vast majority of the reviewed biomass equations (127 in total) took the simple linear form

$$\text{Log}(M) = A + B \times \text{Log}(D) \quad (1)$$

where Log(M) is either the natural or the 10-base logarithmic transformation of the biomass data for different tree components, Log(D) is the diameter at breast height (either in natural or 10-base logarithmic transformation) and A and B the estimated parameters. In 200 regressions tree height was entered as the second independent variable or was used in combination with D. In the 280 empirical regressions, D was the only independent variable and the mathematical relationship between tree biomass and D fell into several formulae (see Appendix A).

The compiled equations do not refer to the same spatial scale; some of them were built on data obtained from a single stand, whereas others (e.g. Marklund's (1987, 1988) equations for the main tree species of Sweden) are based on data from large geographical areas. There are no such equations for temperate or Mediterranean conditions. The amount of sampled trees varied from 3 to 1503 with between 6 and 40 as the most usual amount (Fig. 1a). Only Marklund's (1987, 1988) equations are consistently based on a sample size of several hundred felled trees. In 175 equations


**Table 2.** Number of compiled biomass equations according to tree species and tree component. For the abbreviations see Table 1.

	AB	ABW	BR	CO	CR	DB	FL	FL(i)	RC	RF	RS	RT	SB	SR	ST	SU	SW	TB	TW	Total
<i>Abies balsamea</i>	—	—	—	—	—	—	—	—	—	—	—	—	4	—	—	—	—	—	—	4
<i>Abies</i> spp.	—	—	—	—	2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2
<i>Acer pseudoplatanus</i>	—	2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2
<i>Alnus glutinosa</i>	2	1	3	—	—	—	2	—	—	—	—	—	—	3	—	—	—	—	—	11
<i>Alnus incana</i>	2	—	2	—	—	—	2	—	—	—	—	—	—	2	—	—	—	—	—	8
<i>Arbutus unedo</i>	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3
<i>Betula pendula</i>	1	1	2	—	—	—	1	—	—	—	—	2	—	2	—	—	—	—	—	9
<i>Betula pubescens</i>	1	—	1	—	—	—	1	—	—	—	—	2	—	1	—	2	—	—	—	8
<i>Betula pubescens</i> ssp. <i>czerpanovii</i>	—	—	1	—	—	1	1	—	—	—	—	—	1	—	—	—	—	—	—	4
<i>Betula</i> spp.	—	4	3	—	2	4	2	—	—	—	—	1	4	—	1	1	5	—	—	27
<i>Eucalyptus</i> spp.	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1
<i>Fagus crenata</i>	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1
<i>Fagus moesiacaca</i>	1	—	1	—	—	—	1	—	—	—	—	—	—	1	1	—	—	—	—	5
<i>Fagus sylvatica</i>	8	4	7	—	4	—	6	—	1	1	1	4	2	—	8	—	2	—	—	48
<i>Fraxinus excelsior</i>	—	2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2
<i>Larix sibirica</i>	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	2
<i>Larix</i> spp.	—	—	—	—	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1
<i>Picea abies</i>	16	1	27	—	17	13	28	—	2	—	2	7	14	1	16	1	12	3	3	159
<i>Picea engelmannii</i>	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	2
<i>Picea rubens</i>	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1
<i>Picea sitchensis</i>	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1
<i>Picea</i> spp.	1	—	—	—	3	—	—	—	—	—	—	—	—	1	—	—	—	—	—	5
<i>Pinus banksiana</i>	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1
<i>Pinus contorta</i>	1	—	—	—	1	—	—	—	—	—	3	—	—	1	—	—	—	—	—	6
<i>Pinus nigra</i> var. <i>maritima</i>	—	—	—	—	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—	2
<i>Pinus pinaster</i>	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1
<i>Pinus radiata</i>	1	—	—	—	—	—	—	—	—	—	2	—	—	—	—	—	—	—	—	3
<i>Pinus sylvestris</i>	27	4	26	3	11	12	32	2	7	1	1	7	15	3	23	2	13	—	—	191
<i>Pinus taeda</i>	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1
<i>Populus tremula</i>	2	—	1	—	1	—	1	—	—	—	—	—	—	2	—	—	—	—	—	7
<i>Populus trichocarpa</i>	1	—	—	—	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—	3
<i>Pseudotsuga menziesii</i>	3	1	1	—	2	—	1	—	—	—	6	—	—	1	—	—	—	—	—	15
<i>Pseudotsuga</i> spp.	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1
<i>Quercus conferta</i>	—	2	8	—	—	—	—	—	—	—	—	—	—	—	6	—	—	—	—	16
<i>Quercus ilex</i>	10	1	8	—	1	—	6	—	3	—	3	4	—	6	—	—	—	—	—	42
<i>Quercus petraea</i>	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1
<i>Quercus pyrenaica</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	1
<i>Quercus</i> spp.	1	4	—	—	2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	7
<i>Tilia cordata</i>	—	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Total	83	29	91	3	50	30	84	2	13	2	7	48	37	4	72	5	41	3	3	607

**Table 3.** Geographical distribution of the compiled biomass equations. The numbers indicate the total number of equations for all tree components and for each country. Studies for which the region was not specified are indicated by n/a.

	A	B	CZ	DK	FI	FR	DE	GR	IC	I	NL	N	PL	P	SP	SE	UK	Eur	n/a	Total
<i>Abies balsamea</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4	4	
<i>Abies spp.</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	—	—	2	
<i>Acer pseudoplatanus</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	—	—	2	
<i>Alnus glutinosa</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	8	3	—	—	11	
<i>Alnus incana</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	8	—	—	—	8	
<i>Arbutus unedo</i>	—	—	—	—	—	—	—	—	—	3	—	—	—	—	—	—	—	—	3	
<i>Betula pendula</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4	3	—	2	9	
<i>Betula pubescens</i>	—	—	—	—	4	—	—	—	—	—	—	—	—	—	4	—	—	—	8	
<i>Betula pubescens</i> ssp. <i>czerepanovii</i>	—	—	—	—	4	—	—	—	—	—	—	—	—	—	—	—	—	—	4	
<i>Betula spp.</i>	—	—	—	—	—	9	—	—	—	—	—	—	—	—	14	4	—	—	27	
<i>Eucalyptus spp.</i>	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1	
<i>Fagus crenata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	
<i>Fagus moesiaca</i>	—	—	—	—	—	—	—	5	—	—	—	—	—	—	—	—	—	—	5	
<i>Fagus sylvatica</i>	1	2	3	—	—	9	2	—	—	10	10	—	—	4	5	2	—	—	48	
<i>Fraxinus excelsior</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	—	—	2	
<i>Larix sibirica</i>	—	—	—	—	—	—	—	—	2	—	—	—	—	—	—	—	—	—	2	
<i>Larix spp.</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1	
<i>Picea abies</i>	4	4	16	7	21	—	54	—	4	—	—	12	—	—	36	1	5	—	159	
<i>Picea engelmannii</i>	—	—	—	—	—	—	—	—	2	—	—	—	—	—	—	—	—	—	2	
<i>Picea rubens</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	1	
<i>Picea sitchensis</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1	
<i>Picea spp.</i>	—	—	—	—	—	—	—	—	3	—	—	—	—	—	—	2	—	—	5	
<i>Pinus banksiana</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	1	
<i>Pinus contorta</i>	—	—	—	—	—	—	—	—	2	—	—	—	—	—	—	2	—	2	6	
<i>Pinus nigra</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	—	—	2	
var <i>maritima</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	—	—	2	
<i>Pinus pinaster</i>	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1	
<i>Pinus radiata</i>	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	2	3	
<i>Pinus sylvestris</i>	—	6	49	—	44	—	—	—	—	—	27	17	—	—	27	13	3	3	191	
<i>Pinus taeda</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	1	
<i>Populus tremula</i>	—	—	—	—	—	—	3	—	—	—	—	—	—	—	4	—	—	—	7	
<i>Populus trichocarpa</i>	—	—	—	—	—	—	—	—	3	—	—	—	—	—	—	—	—	—	3	
<i>Pseudotsuga menziesii</i>	—	—	—	—	—	—	—	—	—	1	7	—	—	—	—	2	—	5	15	
<i>Pseudotsuga spp.</i>	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1	
<i>Quercus conferta</i>	—	—	—	—	—	—	—	16	—	—	—	—	—	—	—	—	—	—	16	
<i>Quercus ilex</i>	—	—	—	—	—	—	—	—	5	—	—	—	—	37	—	—	—	—	42	
<i>Quercus petraea</i>	—	—	—	—	—	1	—	—	—	—	—	—	—	—	—	—	—	—	1	
<i>Quercus pyrenaica</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	1	
<i>Quercus spp.</i>	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6	—	—	7	
<i>Tilia cordata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1	
Total	6	12	66	7	80	10	59	21	16	22	18	39	17	1	41	109	49	8	22	607

A=Austria, B=Belgium, CZ=Czech republic, DK=Denmark, FI=Finland, FR=France, DE=Germany, GR=Greece, IC=Iceland, I=Italy, NL=Netherlands, N=Norway, PL=Poland, P=Portugal, SP=Spain, SE=Sweden, UK= United Kingdom, Eur=Europe




**Fig. 1.** Frequencies of a) the biomass and b) the volume equations according to the number of sampled trees used for the development of the equation.

the number of sampled trees upon which the estimation of the empirical parametric values had been based was not reported.

The range of size of the sampled trees varied for each equation (Appendix A), implying that diameter and height range should be taken into account when applicability of the equations is evaluated. Our analysis also indicated that different equations generate different biomass predictions for trees of the same size (Fig. 2). The difference between predicted values of foliage biomass was large, whereas the predicted total aboveground biomass values of *Picea abies* was

relatively consistent (Figs. 2a–b). The number of biomass equations available for roots was small and the differences between predicted root biomass values were high (Fig 2c).

The value of the coefficient of determination ( $r^2$ ) was reported in most of the regressions and varied from 0.012 to 0.99. Especially, the biomass of dead branches of Norway spruce seemed to be difficult to estimate accurately. In general, equations with notably low  $r^2$ -values are excluded, but those obtained for dead branched were kept to show overall difficulties in prediction of the biomass of this component. Only in about 1/10



**Fig. 2.** Predicted foliage biomass a) and total aboveground biomass of *Picea abies* b), and root biomass of *Pinus sylvestris* c) as a function of tree diameters (D). The biomass equations were retrieved from Appendix A. The range of diameter of the illustrated equations indicates the range of observations in the original data on which the equation is based on. When the range of original observation was not reported a minimum of 10 cm and a maximum of 40 cm for diameter was used in this figure.

of the papers concerning biomass equations are some kind of error estimates for the equations presented. The forms of the error estimates are diverse and vary from article to article.

### 3.2. Stem Volume Equations


The total number of the compiled stem volume equations was 230 (Appendix B), and they covered 55 tree species altogether (Table 4). Most of the European countries have already developed

stem volume equations mainly for the planning of the use of forest resources. However, there is no straightforward, commonly accepted definition for stem volume in Europe. In general, the volume of stemwood extending from root collar up to the top of the stems is accounted in the equations developed in the Nordic countries. For some of the reviewed regressions, the stem is the part of the main trunk up to a minimum diameter of 7 cm (it is usually called the merchantable volume) while some authors have not reported definition of the stem related to their equations.


**Table 4.** Geographical distribution of the compiled stem volume equations. The numbers indicate the total number of equations for each country.

Scientific name	A	B	CR	CZ	FI	DE	IC	I	NL	N	PL	R	SE	UK	Total
<i>Abies alba</i>	—	—	—	—	—	—	—	—	—	1	—	—	—	—	1
<i>Abies grandis</i>	—	—	—	—	—	—	—	—	1	1	—	—	—	—	2
<i>Abies sibirica</i>	—	—	—	—	—	1	—	—	—	—	—	—	—	—	1
<i>Abies spp.</i>	2	—	—	—	—	—	—	—	—	—	—	1	—	—	3
<i>Acacia spp.</i>	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1
<i>Acer pseudoplatanus</i>	—	1	—	—	—	—	—	—	1	—	—	1	—	1	4
<i>Alnus alba</i>	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1
<i>Alnus glutinosa</i>	—	—	—	—	—	—	—	—	1	2	—	—	3	—	6
<i>Alnus incana</i>	—	—	—	—	—	—	—	—	—	1	—	—	—	—	1
<i>Alnus nigra</i>	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1
<i>Alnus spp.</i>	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
<i>Arbutus unedo</i>	—	—	—	—	—	—	—	1	—	—	—	—	—	—	1
<i>Betula pendula</i>	—	—	—	—	—	—	—	—	1	—	—	—	—	—	1
<i>Betula spp.</i>	—	1	—	—	4	—	—	—	—	2	—	1	6	1	15
<i>Carpinus spp.</i>	—	—	—	—	—	—	—	—	3	—	—	—	—	—	3
<i>Chamaecyparis lawsoniana</i>	—	—	—	—	—	—	—	—	1	—	—	—	—	—	1
<i>Corylus avellana</i>	—	—	—	—	—	—	—	—	—	1	—	—	—	—	1
<i>Fagus spp.</i>	2	—	—	—	—	—	—	—	—	—	—	1	—	1	4
<i>Fagus sylvatica</i>	—	1	—	—	—	2	—	—	2	—	—	—	—	—	5
<i>Fraxinus excelsior</i>	—	1	—	—	—	—	—	—	1	—	—	—	5	—	7
<i>Fraxinus spp.</i>	—	—	—	—	—	—	—	—	—	1	—	1	—	1	3
<i>Larix decidua</i>	2	1	—	—	—	—	—	—	1	1	—	—	—	—	5
<i>Larix hybrid</i>	—	—	—	—	—	—	—	—	—	1	—	—	—	—	1
<i>Larix kaempferi</i>	—	—	—	—	—	—	—	—	1	1	—	—	—	—	2
<i>Larix sibirica</i>	—	—	—	—	—	—	3	—	—	1	—	—	—	—	4
<i>Larix spp.</i>	—	—	—	—	—	2	—	—	3	—	—	1	—	—	6
<i>Picea abies</i>	2	1	—	2	7	2	1	—	5	12	2	—	8	—	42
<i>Picea engelmannii</i>	—	—	—	—	—	—	1	—	—	—	—	—	—	—	1
<i>Picea sitchensis</i>	—	—	—	—	—	—	—	—	1	3	—	—	—	—	4
<i>Picea spp.</i>	—	—	—	—	—	—	1	—	—	—	—	1	—	—	2
<i>Pinus contorta</i>	—	—	—	—	—	—	1	—	1	—	—	—	3	—	5
<i>Pinus nigra var maritima</i>	—	—	—	—	—	—	—	—	1	—	—	—	—	—	1
<i>Pinus nigra var nigra</i>	—	—	—	—	—	—	—	—	1	—	—	1	—	—	2
<i>Pinus spp.</i>	—	—	—	—	—	2	—	—	3	—	—	—	—	—	5
<i>Pinus sylvestris</i>	1	2	—	—	8	1	—	1	4	8	—	1	8	—	34
<i>Populus spp.</i>	1	—	—	—	—	—	—	—	3	—	—	2	—	1	7
<i>Populus tremula</i>	—	—	—	—	—	—	—	—	—	2	—	1	3	—	6
<i>Populus trichocarpa</i>	—	—	—	—	—	—	1	—	—	—	—	—	—	—	1
<i>Prunus avium</i>	—	1	—	—	—	—	—	—	—	—	—	—	—	—	1
<i>Pseudotsuga menziesii</i>	—	1	—	—	—	—	—	—	1	1	—	1	—	—	4
<i>Pseudotsuga spp.</i>	—	—	—	—	—	—	—	—	3	—	—	—	—	—	3
<i>Quercus grisea</i>	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1
<i>Quercus ilex</i>	—	—	1	—	—	—	—	1	—	—	—	—	—	—	2
<i>Quercus laevis</i>	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1
<i>Quercus pubescens</i>	—	—	2	—	—	—	—	—	—	—	—	1	—	—	3
<i>Quercus robur</i>	—	—	—	—	—	—	—	—	1	—	—	—	—	—	1
<i>Quercus rubra</i>	—	1	—	—	—	—	—	—	1	—	—	—	—	—	2
<i>Quercus spp.</i>	2	1	—	—	—	—	—	—	3	—	—	1	—	1	8
<i>Salix caprea</i>	—	—	—	—	—	—	—	—	—	1	—	1	—	—	2
<i>Salix spp.</i>	—	—	—	—	—	—	—	—	—	—	—	2	—	—	2
<i>Sorbus aucuparia</i>	—	—	—	—	—	—	—	—	—	1	—	—	—	—	1
<i>Thuja pilicata</i>	—	—	—	—	—	—	—	—	1	1	—	—	—	—	2
<i>Tilia cordata</i>	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1
<i>Tsuga heterophylla</i>	—	—	—	—	—	—	—	—	1	1	—	—	—	—	2
<i>Ulmus spp.</i>	—	1	—	—	—	—	—	—	1	—	—	1	—	—	3
Total	13	13	3	2	21	8	8	3	47	43	2	25	36	6	230

A=Austria, B=Belgium, CR=Croatia, CZ=Czech republic, FI=Finland, DE=Germany, IC=Iceland, I=Italy, NL=Netherlands, N=Norway, PL=Poland, R=Romania, SE=Sweden, UK= United Kingdom



**Fig. 3.** Predicted stem volume of *Fagus sylvatica* as a function of tree diameter ( $D$ ). The volume equations are presented in Appendix B.



**Fig. 4.** Predicted stem volume of *Pinus sylvestris* as a function of tree diameter ( $D$ ). The volume equations are presented in Appendix B.

If the definition of the stem volume is reported in the original paper, it is indicated in the comment field of the appendix table.

The major part of the stem volume regressions was for coniferous tree species (Table 4). Thirty-nine equations were reported for Norway spruce (*Picea abies* (L.) Karst.) and 31 for Scots pine (*Pinus sylvestris* L.), from which more than half were built for Scandinavian countries. For the broadleaved tree species within the genera of *Betula*, *Fagus*, and *Quercus* the number of available equations were 16, 9, and 18, respectively.

Most of the stem volume equations were based on a sample size of several hundred or a few thousand felled trees (Fig. 1b). Only three equations were based on a sample size of more than 5000 trees. In 97 of the equations the number of sample trees was not reported.

In almost every of the compiled stem volume equations the independent variables were  $D$  and/or  $H$  or various mathematical combinations of these. However, in three equations the formula used to fit the tree-scale data was solely based on  $D$  and was a simple power function (simple allometric equation). More than two parameters were incorporated into the formulae of 212 stem volume regressions, and 18 equations included six parameters. The number of the sampled trees

(from which the empirical stem volume regressions were obtained) varied from five to more than 7446 (Appendix B). The range of diameters of the sampled trees varied between equations and for 125 of the compiled stem volume equations the range was not reported. In all the compiled equations the coefficient of determination was more than 0.58 irrespective of species, location,  $D$  range, site conditions, etc.

Predicted stem volume estimates varied according to the applied equation (Fig. 3 and Fig. 4). For example, the stem volume of a beech tree with a diameter of 40 cm varies between 1.1 and 2.2  $m^3$  (Fig. 3). On the other hand, all stem volume equations of e.g. Scots pine produced relatively consistent stem volume estimates. The equation reported by Schelhaas et al. (2002) and one of the equations published by Laasasenaho (1982) seemed to deviate from the others (Fig. 4). However, Laasasenaho (1982) reported two other equations which had different form or more explanatory variables (height in addition to dbh), and they gave consistent predictions with the models of other authors.

## 4 Discussion

---

Reliable methods of estimating forest biomass and carbon stocks as well as volume of the growing stock at different spatial and temporal scales and for different biomes are needed. In national forest inventories, emphasis has been placed on the assessment of merchantable timber, and inventories provide highly accurate estimates of the growing stock (Laitat et al. 2000). The current need to assess changes in the forest carbon has arisen as a result of the Climate Convention and the Kyoto Protocol. In general, assessment of forest biomass and carbon stock is based on information on forest resources i.e. estimates of forested area and volume of the growing stock as reported by national forest inventories (Liski and Kauppi 2000). Reported volume estimates are multiplied with simple biomass expansion factors and/or conversion factors to obtain biomass estimates.

In national inventories, the volume of the growing stock is estimated with the help of volume equations. The results of this study show that representative volume equations are available for major tree species in Europe, since volume equations are developed for different vegetation zones and most of the equations are based on a relatively high number of sampled trees. However, the volume equations vary in terms of the dimensions accounted for (merchantable stem volume only or unmerchantable included), and the estimates obtained with different equations cannot be compared or aggregated, and they cannot be converted to biomass estimates by just using a single biomass expansion value. The differences were the most evident with tree species that had irregular branching patterns (e.g. beech), whereas volume equations of e.g. Scots pine were more consistent. The inconsistency of the different volume equations applied to national forest inventories was also reported by Köhl et al. (1997). As national estimates of the volume of the growing stock are converted to biomass estimates, the applicability of the biomass expansion factors to the applied

volume equation needs to be evaluated to avoid highly biased biomass estimates.

Reliability of the national carbon inventories can be improved by applying biomass equations directly to tree-scale measurements of diameter ( $D$ ) at sampled plots of forest inventories (Jalkanen et al. 2005). Consequently, the additional source of error introduced by conversion or expansion factors can be avoided. The compiled database on biomass equations provides a basis for the selection of the applicable biomass equation when representative national equations are not available. The database can be also used as a source of reference for the development of local equations. Since the number of sampled trees used for the development of the biomass equations seemed to be relatively small, it is necessary to use several equations rather than only one in order to obtain unbiased predictions.

The analysis of the collected information showed that both species coverage and the spatial distribution of the equations is limited. The vast majority of the models were developed for coniferous tree species growing in northern and central European forest ecosystems. Only a small number of biomass and stem volume regressions were collected for tree species in the eastern and southern parts of Europe (Tables 3 and 4). A rather limited number of equations for the estimation of root biomass has been compiled so far, indicating that a more extensive survey should take place and that more root biomass datasets should be collected across Europe. In a similar study conducted in Australia, Snowdon et al. (2000) stressed that more root biomass studies are needed and suggested that fractal geometry could be a promising tool to overcome the practical problems arising from the destructive sampling of belowground tree biomass. Ter-Mikaelian and Korzukhin (1997) reported no equations for estimating the root biomass of tree species growing in the USA.

Most of the collected equations lack information

on the error estimates of the empirical parameters. According to Keith et al. (2000), the main sources of error in implementing allometric regressions could occur at the treescale and when biomass estimates are extrapolated from plot to regional scale (see also Satoo and Madgwick 1982). It should be noticed that when a logarithmic or any other transformation is applied to the raw data, biomass and stem volume predictions are biased (Baskerville 1972, Sprugel 1983). Mathematical formulae for correcting bias provide accurate estimates even though assumptions about the distribution of statistical errors must be made. The inherent bias arising from data transformation could be eliminated if iterative procedures were to be applied to the data (for a more detailed discussion see Payandeh 1981).

Biased predictions may also be obtained when the sum of biomass estimates (developed for different tree components i.e., stem, crown and roots) does not match the predictions derived from the total biomass equation (what is called the additivity problem). Parresol (2001) provided statistical methods to account for this bias while Snowdon et al. (2000) reported that the additivity problem does not appear when allometric equations are developed from non-transformed data. Another statistical problem is caused by collinearity or multicollinearity, where the independent variables in a regression analysis are themselves correlated (Ott 1993). Thus, the value of the coefficient of determination in stem volume and biomass equations (with more than one independent variable) may not be a reliable criterion for the choice of the best-fitting equation, and biased predictions may be obtained when this problem is not taken into account. However, the collinearity problem is seldom mentioned in original papers, where more often than not, diameter and height are the independent variables in estimating either stem volume or tree biomass.

The equations presented in this review can be used for national biomass and carbon inventories, for ecological studies, for validating theoretical models and for planning the use of forest resources. Since the original biomass studies may have been conducted for very specific purposes, following different sampling procedures and perhaps atypical stand structures, the applicability of an equation to its intended purpose needs to be evaluated in terms of the geographical distribution of the sampled population, the number of sampled trees, the range of dimensions ( $D, H$ ) of sampled trees, accounted dimensions and applied definitions.

Pooled equations based on raw data collected from wide geographical areas may also provide a promising alternative to estimate biomass changes at the landscape scale Wirth et al. 2004). The empirical models reviewed in this article may also be used in order to build generalised stem volume and biomass equations for different species and different tree components (see Pastor et al. 1983/1984 for American species and Zianis and Mencuccini 2003 for the genus *Fagus*), to develop BEF for tree species across Europe (Lehtonen et al. 2004) and to validate process-based models of forest productivity.

## References

---

- Baskerville, G.L. 1972. Use of logarithmic regression in the estimation of plant biomass. *Canadian Journal of Forestry* 2: 49–53.
- Brown, S. 1997. Estimating biomass and biomass change of tropical forests, a premier. FAO Forestry Paper 134.
- Bruce, D. & Schumacher, F.X. 1950. Forest mensuration. McGraw-Hill Book Company, Inc. New York. 483 p.
- Clark, D.A., Brown, S., Kicklighter, D.W., Chambers, J.Q., R, T.J. & Ni, J. 2001. Measuring net primary production in forests: concepts and field methods. *Ecological Applications* 11(2): 356–370.
- Drake, J.B., Dubayah, R.O., Knox, R.G., Clark, D.B. & Blair, J.B. 2002. Sensitivity of large-footprint lidar to canopy structure and biomass in a neotropical rainforest. *Remote Sensing of Environment* 81: 378–392.
- Eamus, D., McGuinness, K. & Burrows, W. 2000. Review of allometric relationships for estimating woody biomass for Queensland, the Northern Territory and Western Australia. National Carbon Accounting System Technical Report. Series Review of allometric relationships for estimating woody biomass for Queensland, the Northern Territory and Western Australia 5A. Australian Greenhouse Office, Canberra. 56 p.
- Hall, D.O. 1997. Biomass energy in industrialised countries – a view of the future. *Forest Ecology and Management* 91: 17–45.
- IPCC, 2003. Report on good practice guidance for land use, land-use change and forestry. IPCC National Greenhouse Gas Inventories Programme <http://www.ipcc-nrgip.iges.or.jp/public/gpglulucf/gpglulucf.htm>, Japan.
- Jalkanen, A., Mäkipää, R., Ståhl, G., Lehtonen, A. & Petersson, H. 2005. Estimation of biomass stock of trees in Sweden: comparison of biomass equations and age-dependent biomass expansion factors. *Annals of Forest Science* 62 (In press)
- Jenkins, J.C., Chojnacky, D.C., Heath, L.S. & Birdsey, R.A. 2003. National-scale biomass estimators for United States tree species. *Forest Science* 49: 12–35.
- , Chojnacky, D.C., Heath, L.S. & Birdsey, R.A. 2004. Comprehensive database of diameter-based biomass regressions for North American tree species. Gen. Tech. Rep. NE-319. US Forest Service. 45 p.
- Keith, H., Barrett, D. & Keenan, R. 2000. Review of allometric relationships for estimating woody biomass for New South Wales, the Australian Capital Territory, Victoria, Tasmania, and South Australia. National Carbon Accounting System Technical Report. Series Review of allometric relationships for estimating woody biomass for New South Wales, the Australian Capital Territory, Victoria, Tasmania, and South Australia 5B. Australian Greenhouse Office, Canberra. 114 p.
- Köhl, M., Päivinen, R., Traub, B. & Miina, S. 1997. Comparative study. In: Study on European forestry information and communication system. Reports on forest inventory and survey systems 2. European Commission. p. 1265–1322.
- Laitat, E., Karjalainen, T., Loustau, D. & Lindner, M. 2000. Towards an integrated scientific approach for carbon accounting in forestry. *Biotechnology, Agronomy, Society and Environment* 4: 241–251.
- Lehtonen, A., Mäkipää, R., Heikkinen, J., Sievänen, R. & Liski, J. 2004. Biomass expansion factors (BEFs) for Scots pine, Norway spruce and birch according to stand age for boreal forests. *Forest Ecology and Management* 188: 211–224.
- Liski, J. & Kauppi, P. 2000. Carbon cycle and biomass. In: FAO (ed.). Forest resources of Europe, CIS, North America, Australia, Japan and New Zealand (industrialized temperate/boreal countries). UN-ECE/FAO Contribution to the Global Forest Resources Assessment 2000, Main Report (in press). United Nations, New York and Geneva. p. 155–171.
- Löwe, H., Seufert, G. & Raes, F. 2000. Comparison of methods used within member states for estimating

- CO<sub>2</sub> emissions and sinks to UNFCCC and UE monitoring mechanism: forest and other wooded land. *Biotechnology, Agronomy, Society and Environment* 4: 315–319.
- Marklund, L.G. 1987. Biomass functions for Norway spruce (*Picea abies* (L.) Karst.) in Sweden. Sveriges lantbruksuniversitet, Institutionen för skogstaxering, Rapport 43. 127 p.
- 1988. Biomassfunktioner för tall, gran och björk i Sverige. Sveriges lantbruksuniversitet, Institutionen för skogstaxering, Rapport 45. 73 p.
- Montes, N., Gauquelin, T., Badri, W., Bertaudiere, V. & Zaoui, E.H. 2000. A non-destructive method for estimating above-ground forest biomass in threatened woodlands. *Forest Ecology and Management* 130: 37–46.
- Ott, R.L. 1993. An introduction to statistical methods and data analysis. Duxbury press, California. 132 p.
- Parresol, R.B. 2001. Additivity of nonlinear biomass equations. *Canadian Journal of Forest Research* 31: 865–878.
- Pastor, J., Aber, J.D. & Melillo, J.M. 1983/1984. Biomass prediction using generalized allometric regressions for some northeast tree species. *Forest Ecology and Management* 7: 265–274.
- Payandeh, B. 1981. Choosing regression models for biomass prediction equations. *The Forestry Chronicle* 57: 230–232.
- Santantonio, D., Hermann, R.K. & Overton, W.S. 1977. Root biomass studies in forest ecosystems. *Pedobiologia* 17: 1–31.
- Satoo, T. & Madgwick, H.A.I. 1982. Forest biomass. Kluwer Academic Publishers Group, London. 160 p.
- Snowdon, P., Eamus, D., Gibbons, P., Khanna, P.K., Keith, H., Raison, R.J. & Kirschbaum, M.U.F. 2000. Synthesis of allometrics, review of root biomass and design of future woody biomass sampling strategies. National Carbon Accounting System Technical Report. Series Synthesis of allometrics, review of root biomass and design of future woody biomass sampling strategies 17. Australian Greenhouse Office, Canberra. 114 p.
- Sprugel, D.G. 1983. Correcting for bias in log-transformed allometric equations. *Ecology* 64: 209–210.
- Ter-Mikaelian, M.T. & Korzukhin, M.D. 1997. Biomass equations for sixty-five North American tree species. *Forest Ecology and Management* 97: 1–24.
- Tritton, L.M. & Hornbeck, J.W. 1982. Biomass equations for major tree species of the Northeast. Series Biomass equations for major tree species of the Northeast NE-69. U.S. Department of Agriculture, Northeastern Forest Experiment Station, General Technical Report. 46 p.
- Wirth, C., Schumacher, J. & Schulze, E.-D. 2004. Generic biomass functions for Norway spruce in Central Europe – a meta-analysis approach toward prediction and uncertainty estimation. *Tree Physiology* 24: 121–139.
- Zianis, D. & Mencuccini, M. 2003. Aboveground biomass relationship for beech (*Fagus moesiaca* Cz.) trees in Vermio Mountain, Northern Greece, and generalised equations for *Fagus* spp. *Annals of Forest Science* 60: 439–448.
- 2004. On simplifying allometric analyses of forest biomass. *Forest Ecology and Management* 187: 311–332.

*Total of 33 references*

**Appendix A.** Biomass equations for different biomass components by tree species (see abbreviations for dependent variable from table 1). In addition to scientific names of the tree species, common names are shown as they are reported in the original publications. The format of the biomass equation is given in the column labelled Equation, and a, b, c, d, and e are parameter values. The “ln” is the natural logarithm and the “log” is the



		Unit of Biom.	D	H	D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	r <sup>2</sup>
<i>Abies balsamea</i>											
1 –	log(RT)	kg	cm	–	–	–	68	8	–	–	–
2 –	log(RT)	kg	cm	–	–	–	68	8	89	0.92	
3 –	log(RT)	kg	cm	–	–	–	68	8	40	0.928	
4 –	log(RT)	kg	cm	–	–	–	68	8	40	0.898	
<i>Abies</i> spp. (Fir)											
5 UK	CR	t	cm	–	–	–	9	6	1	–	–
6 UK	CR	t	cm	–	–	–	9	6	2	–	–
<i>Acer pseudoplatanus</i> (Sycamore)											
7 UK	ln(ABW)	kg	cm	–	3.7–31	–	10	14	10	0.991	
8 UK	ln(ABW)	kg	cm	–	3.5–28	–	10	14	15	0.995	
<i>Alnus glutinosa</i> (Common alder, Black alder, Klubbal)											
9 Sweden	AB	kg	mm	–	1–17.3	2.5–17.6	39	4	–	0.987	
10 Sweden	AB	kg	mm	–	12.2–28.3	13–25.4	38	4	–	0.98	
11 UK	ABW	kg	cm	–	–	–	32	14	12	0.985	
12 Sweden	BR	kg	mm	–	12.2–28.3	13–25.4	38	4	–	0.66	
13 Sweden	BR	kg	mm	–	1–17.3	2.5–17.6	39	4	–	0.922	
14 UK	BR	kg	cm	–	–	–	32	14	12	0.924	
15 Sweden	FL	kg	mm	–	12.2–28.3	13–25.4	38	4	–	0.47	
16 Sweden	FL	kg	mm	–	1–17.3	2.5–17.6	39	4	–	0.927	
17 Sweden	ST	kg	mm	–	12.2–28.3	13–25.4	38	4	–	0.82	
18 Sweden	ST	kg	mm	–	1–17.3	2.5–17.6	39	4	–	0.969	
19 UK	ST	kg	cm	–	–	–	32	14	12	0.991	
<i>Alnus incana</i> (Grey alder, Gråal, Harmaaleppä)											
20 Sweden	AB	kg	mm	–	0.7–9.3	2–14.8	39	4	–	0.983	
21 Sweden	AB	kg	mm	–	8.9–24.6	13–25.3	38	4	–	0.92	
22 Sweden	BR	kg	mm	–	0.7–9.3	2–14.8	39	4	–	0.862	
23 Sweden	BR	kg	mm	–	8.9–24.6	13–25.3	38	4	–	0.6	
24 Sweden	FL	kg	mm	–	0.7–9.3	2–14.8	39	4	–	0.64	
25 Sweden	FL	kg	mm	–	8.9–24.6	13–25.3	38	4	–	0.44	
26 Sweden	ST	kg	mm	–	0.7–9.3	2–14.8	39	4	–	0.98	
27 Sweden	ST	kg	mm	–	8.9–24.6	13–25.3	38	4	–	0.89	
<i>Arbutus unedo</i> (Strawberry-tree)											
28 Italy	AB	kg	cm	–	6–15	–	7	8	26	0.955	
29 Italy	ABW	kg	cm	–	6–15	–	7	8	3	26	0.955
30 Italy	CR	kg	cm	–	6–15	–	7	8	26	0.955	
<i>Betula pendula</i> (Silver birch, Pendula birch, White birch, Rauduskoivu, Värtnbjörk)											
31 Sweden	AB	kg	mm	–	1.8–13.7	3.2–19.9	35	4	–	0.985	
32 UK	ABW	kg	cm	–	–	–	32	14	13	0.99	
33 Sweden	BR	kg	mm	–	1.8–13.7	3.2–19.9	35	4	–	0.747	
34 UK	BR	kg	cm	–	–	–	32	14	13	0.99	
35 Sweden	FL	kg	mm	–	1.8–13.7	3.2–19.9	35	4	–	0.884	
36 –	log(RT)	kg	cm	–	–	–	68	8	3	0.983	
37 –	log(RT)	kg	cm	m	–	–	68	8	3	0.997	
38 Sweden	ST	kg	mm	–	1.8–13.7	3.2–19.9	35	4	–	0.979	
39 UK	ST	kg	cm	–	–	–	32	14	13	0.99	
<i>Betula pubescens</i> (White birch, Pubescent birch, Hieskoivu, Glasbjörk, Björk)											
40 Sweden	AB	kg	mm	–	0.8–8.5	2.3–12	35	4	–	0.977	
41 Sweden	BR	kg	mm	–	0.8–8.5	2.3–12	35	4	–	0.875	
42 Sweden	FL	kg	mm	–	0.8–8.5	2.3–12	35	4	–	0.918	
43 Finland	SB	kg	cm	m	2–16	4.6–16.7	58	8	53	0.986	
44 Finland	SB	kg	cm	m	1.3–13	3.3–13.2	58	8	56	0.984	
45 Sweden	ST	kg	mm	–	0.8–8.5	2.3–12	35	4	–	0.966	

10-based logarithm. Number of sampled trees ( $n$ ), coefficients of determination ( $r^2$ ), and range of diameter ( $D$ ) and height ( $H$ ) of sampled trees are reported when available in the original article. References (Ref.) to the original papers according to author as well as the contact (Cont.) person who submitted the equation to this database are given at the end of the table. In the comments column (Comm.) occur some comments about the particular equation.

Equation	a	b	Parameters		
			c	d	e
1	a-log(D)+b	2.4452	-1.7143	-	-
2	a-log(D)+b	2.45	0.681	-	-
3	a-log(D)+b	2.0027	0.0629	-	-
4	a-log(D)+b	2.4613	-0.4023	-	-
5	a·D <sup>b</sup>	5.2193·10 <sup>-4</sup>	1.459	-	-
6	a+b·D <sup>c</sup>	0.0060722	9.58·10 <sup>-6</sup>	2.5578	-
7	a+b·ln(D)	-2.7606	2.5189	-	-
8	a+b·ln(D)	-2.7018	2.5751	-	-
9	a·D <sup>b</sup>	0.00079	2.28546	-	-
10	a·D <sup>b</sup>	0.003090	2.022126	-	-
11	a·D <sup>b</sup>	0.0859	2.3537	-	-
12	a·D <sup>b</sup>	0.000003	2.880598	-	-
13	a·D <sup>b</sup>	0.0000006	3.28106	-	-
14	a·D <sup>b</sup>	0.0146	2.5191	-	-
15	a·D <sup>b</sup>	0.000003	2.547045	-	-
16	a·D <sup>b</sup>	0.00239	1.32535	-	-
17	a·D <sup>b</sup>	0.005609	1.888345	-	-
18	a·D <sup>b</sup>	0.00119	2.17247	-	-
19	a·D <sup>b</sup>	0.0841	2.4501	-	-
20	a·D <sup>b</sup>	0.00030	2.42847	-	-
21	a·D <sup>b</sup>	0.000499	2.337592	-	-
22	a·D <sup>b</sup>	0.00001	2.65455	-	-
23	a·D <sup>b</sup>	0.000100	2.297058	-	-
24	a·D <sup>b</sup>	0.00001	2.44406	-	-
25	a·D <sup>b</sup>	0.000076	2.02604	-	-
26	a·D <sup>b</sup>	0.00029	2.40128	-	-
27	a·D <sup>b</sup>	0.000368	2.335763	-	-
28	a+b·D <sup>2</sup>	-2.7563	0.3045	-	-
29	a+b·D <sup>2</sup>	-2.8816	0.2639	-	-
30	a+b·D <sup>2</sup>	0.1253	0.040617	-	-
31	a·D <sup>b</sup>	0.00087	2.28639	-	-
32	a·D <sup>b</sup>	0.2511	2.29	-	-
33	a·D <sup>b</sup>	0.00002	2.63001	-	-
34	a·D <sup>b</sup>	0.0742	2.24	-	-
35	a·D <sup>b</sup>	0.00371	1.11993	-	-
36	a-log(D)+b	2.3547	-1.3	-	-
37	a-log(H·D <sup>2</sup> )+b	0.9308	-1.8	-	-
38	a·D <sup>b</sup>	0.00080	2.28244	-	-
39	a·D <sup>b</sup>	0.193	2.25	-	-
40	a·D <sup>b</sup>	0.00029	2.50038	-	-
41	a·D <sup>b</sup>	0.00004	2.52978	-	-
42	a·D <sup>b</sup>	0.00090	1.47663	-	-
43	a+b·ln(D <sup>2</sup> ·H)	-2.1909	0.8808	-	-
44	a+b·ln(D <sup>2</sup> ·H)	-2.0706	0.7942	-	-
45	a·D <sup>b</sup>	0.00020	2.54302	-	-

**App. A**

		Unit of Biom.	D	H	Range of D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	$r^2$
46 Finland	SW	kg	cm	m	2–16	4.6–16.7	58	8		53	0.994
47 Finland	SW	kg	cm	m	1.3–13	3.3–13.2	58	8		56	0.994
<i>Betula pubescens</i> ssp. <i>czerepanovii</i> (Mountain birch)											
49 Finland	ln(BR)	g	mm	–	–	–	72	8		20	0.836
50 Finland	ln(DB)	g	mm	–	–	–	72	8		20	0.622
51 Finland	ln(FL)	g	mm	–	–	–	72	8		20	0.829
48 Finland	ln(ST)	g	mm	–	–	–	72	8		20	0.98
<i>Betula</i> spp. (Birch, Koivu, Björk)											
52 UK	ln(ABW)	kg	cm	–	2.9–30	–	10	14		27	0.985
53 UK	ln(ABW)	kg	cm	–	2.9–26	–	10	14		15	0.984
54 UK	ln(ABW)	kg	cm	–	3.3–16	–	10	14		16	0.984
55 UK	ln(ABW)	kg	cm	–	3.5–23	–	10	14		15	0.987
56 Finland	BR	kg	cm	m	9–28	13–22.4	57	8		20	0.901
57 Sweden	ln(BR)	g	cm	dm	0.9–9.8	1.8–9.2	19	8		66	0.88
58 Sweden	ln(BR)	kg	cm	–	0–35	0–	50	8	4	235	0.924
59 Finland	ln(CR)	kg	mm	–	–	–	29	8		–	0.839
60 Finland	ln(CR)	kg	mm	–	–	–	29	8		–	0.838
61 Finland	DB	kg	cm	m	9.0–28	13–22.4	57	8		20	0.267
62 Sweden	ln(DB)	kg	cm	–	0–35	0–	50	8		212	0.605
63 Sweden	ln(DB)	kg	cm	m	0–35	0–	50	8		212	0.621
64 Sweden	ln(DB)	g	cm	–	0.9–9.8	18–92	19	8		61	0.56
65 Finland	FL	kg	cm	m	9–28	13–22.4	57	8		20	0.906
66 Sweden	ln(FL)	g	cm	dm	0.9–9.8	1.8–9.2	19	8		14	0.92
67 Finland	RT	kg	cm	m	9–28	13–22.4	57	8	5	20	0.994
68 Finland	SB	kg	cm	m	9–28	13–22.4	57	8		20	0.966
69 Sweden	ln(SB)	kg	cm	–	0–35	0–	50	8		212	0.947
70 Sweden	ln(SB)	kg	cm	m	0–35	0–	50	8		212	0.958
71 Sweden	ln(SB)	g	cm	dm	0.9–9.8	1.8–9.2	19	8		66	0.91
72 Sweden	ln(ST)	kg	cm	m	0–35	0–	50	8		240	0.992
73 Finland	SU	kg	cm	m	9–28	13–22.4	57	8		20	0.96
74 Finland	SW	kg	cm	m	9–28	13–22.4	57	8		20	0.99
75 Sweden	ln(SW)	kg	cm	–	0–35	0–	50	8		240	0.982
76 Sweden	ln(SW)	kg	cm	–	0–35	0–	50	8		212	0.97
77 Sweden	ln(SW)	kg	cm	m	0–35	0–	50	8		212	0.99
78 Sweden	ln(SW)	g	cm	dm	0.9–9.8	1.8–9.2	19	8		66	0.99
<i>Eucalyptus</i> spp. (Eucalypt)											
79 Italy	ln(AB)	kg	cm	–	4–25	–	53	14	6	22	0.99
<i>Fagus crenata</i>											
80 –	log(RT)	kg	cm	m	–	–	68	8		7	0.969
<i>Fagus moesiaca</i> (Beech, Oxia)											
81 Greece	ln(AB)	kg	cm	–	5.4–41	9.2–28	76	14		16	0.99
82 Greece	ln(BR)	kg	cm	–	5.4–41	9.2–28	76	14		16	0.97
83 Greece	ln(FL)	kg	cm	–	5.4–41	9.2–28	76	14		16	0.9
84 Greece	ln(ST)	kg	cm	–	5.4–41	9.2–28	76	14		16	0.98
85 Greece	ln(SU)	kg	cm	–	5.4–41	9.2–28	76	14	7	16	0.78
<i>Fagus sylvatica</i> (Beech, European beech, Hêtres, Rotbuche)											
86 Austria	ln(AB)	kg	cm	m	–	–	31	3		42	0.997
87 Belgium	log(AB)	g	cm	–	35–78.8	–	23	12		6	0.995
88 Czech republic	AB	kg	cm	–	5.7–62.1	9.2–33.9	15	2		20	0.974
89 Germany	AB	kg	cm	–	–	–	65	4		–	–
90 Netherlands	AB	kg	cm	m	–	–	5	8		38	0.991
91 Netherlands	AB	kg	cm	–	–	–	5	8		38	0.988
92 Spain	AB	kg	cm	–	4–34.5	6.1–18.4	67	14		7	0.98
93 Sweden	log(AB)	kg	cm	m	12–64	11–29	61	8		–	–
94 Italy	ABW	kg	cm	m	–	–	11	8	8	–	0.993
95 Italy	ABW	kg	cm	m	–	–	11	8	9	–	0.988
96 Italy	ABW	kg	cm	m	–	–	11	8	10	–	0.991
97 Italy	ABW	kg	cm	m	–	–	11	8		–	0.995
98 Belgium	log(BR)	g	cm	–	35–78.8	–	23	12		6	0.981
99 Czech republic	BR	kg	cm	–	5.7–62.1	9.2–33.9	15	2		20	0.806

	Equation	a	b	Parameters	c	d	e
46	a+b·ln(D <sup>2</sup> ·H)	-1.6047	0.9450	—	—	—	—
47	a+b·ln(D <sup>2</sup> ·H)	-1.5195	0.9204	—	—	—	—
48	a+b·ln(D)	-0.305	1.953	—	—	—	—
49	a+b·ln(D)	-3.368	2.041	—	—	—	—
50	a+b·ln(D)	0.525	1.398	—	—	—	—
51	a+b·ln(D)	-0.313	2.140	—	—	—	—
52	a+b·ln(D)	-2.4166	2.4227	—	—	—	—
53	a+b·ln(D)	-2.7584	2.6134	—	—	—	—
54	a+b·ln(D)	-2.1625	2.3078	—	—	—	—
55	a+b·ln(D)	-2.6423	2.4678	—	—	—	—
56	a+b·log(D <sup>2</sup> ·H)	-3.810	1.2911	—	—	—	—
57	a+b·ln(D)+c·H+d·ln[(D <sup>2</sup> )·H]	12.0993	8.5963	0.0406	-2.9662	—	—
58	a+b·[D/(D+10)]	-3.3633	10.2806	—	—	—	—
59	a+b·ln(D)	-10.7699	2.6016	—	—	—	—
60	a+b·ln(D)	-10.2692	2.5124	—	—	—	—
61	a+b·log(D <sup>2</sup> ·H)	-6.510	1.5593	—	—	—	—
62	a+b·[D/(D+5)]	-5.9507	7.9266	—	—	—	—
63	a+b·[D/(D+30)]+c·H+d·ln(H)	-6.6237	11.2872	-0.3081	2.6821	—	—
64	a+b·ln(D)	1.637	1.9554	—	—	—	—
65	a+b·log(D <sup>2</sup> ·H)	-3.454	1.0961	—	—	—	—
66	a+b·ln(D)+c·ln[(D <sup>2</sup> )·H]	10.2953	7.9621	-2.3022	—	—	—
67	a+b·log(D <sup>2</sup> ·H)	-3.887	1.3668	—	—	—	—
68	a+b·log(D <sup>2</sup> ·H)	-2.311	0.9256	—	—	—	—
69	a+b·[D/(D+14)]	-3.2518	10.3876	—	—	—	—
70	a+b·[D/(D+14)]+c·ln(H)	-4.0778	8.3019	0.7433	—	—	—
71	a+b·ln(D)+c·H+d·ln[(D <sup>2</sup> )·H]	5.8227	3.3503	0.0259	-0.8584	—	—
72	a+b·[D/(D+7)]+c·H+d·ln(H)	-3.5686	8.2827	0.0393	0.5772	—	—
73	a+b·log(D <sup>2</sup> ·H)	-3.540	1.1488	—	—	—	—
74	a+b·log(D <sup>2</sup> ·H)	-1.785	0.9910	—	—	—	—
75	a+b·[D/(D+8)]	-3.0932	11.0735	—	—	—	—
76	a+b·[D/(D+11)]	-2.3327	10.8109	—	—	—	—
77	a+b·[D/(D+11)]+c·ln(H)	-3.3045	8.1184	0.9783	—	—	—
78	a+b·ln(D)+c·H+d·ln[(D <sup>2</sup> )·H]	7.4223	3.9941	0.0338	-1.0984	—	—
79	a+b·ln(D)	-1.762	2.2644	—	—	—	—
80	a·log(H·D <sup>2</sup> )+b	0.6816	-1.0003	—	—	—	—
81	a+b·ln(D)	-1.3816	2.3485	—	—	—	—
82	a+b·ln(D)	-5.2898	2.9353	—	—	—	—
83	a+b·ln(D)	-4.1814	1.6645	—	—	—	—
84	a+b·ln(D)	-1.6015	2.3427	—	—	—	—
85	a+b·ln(D)	-1.7716	1.073	—	—	—	—
86	a+b·ln(D)+c·ln(H)	-2.872	2.095	0.678	—	—	—
87	a+b·log(D)	2.85102	2.0666	—	—	—	—
88	a·D <sup>b</sup>	0.453	2.139	—	—	—	—
89	a·D <sup>b</sup>	0.1143	2.503	—	—	—	—
90	a·D <sup>b</sup> ·H <sup>c</sup>	0.0306	2.347	0.590	—	—	—
91	a·D <sup>b</sup>	0.0798	2.601	—	—	—	—
92	a·D <sup>b</sup>	0.1315	2.4321	—	—	—	—
93	a+log[H·(D <sup>2</sup> )]·b	-1.7194	1.0414	—	—	—	—
94	a·D <sup>b</sup> ·H <sup>c</sup>	0.04736	1.80521	0.99603	—	—	—
95	a·D <sup>b</sup> ·H <sup>c</sup>	0.16885	2.44639	-0.1431	—	—	—
96	a·D <sup>b</sup> ·H <sup>c</sup>	0.00868	2.25454	1.09409	—	—	—
97	a·D <sup>b</sup> ·H <sup>c</sup>	0.03927	2.01361	0.87832	—	—	—
98	a+b·log(D)	0.41439	3.18522	—	—	—	—
99	a·D <sup>b</sup>	0.021	2.471	—	—	—	—

## App. A

		Unit of Biom.	D	H		Range of D (cm)	H (m)	Ref.	Cont.	Comm.	n	$r^2$
100 France	ln(BR)	kg	cm	—	—	—	—	42	14	23	0.93	
101 Netherlands	BR	kg	cm	m	—	—	—	5	8	38	0.92	
102 Netherlands	BR	kg	cm	—	—	—	—	5	8	38	0.916	
103 Spain	BR	kg	cm	—	4–34.5	6.1–18.4	—	67	14	7	0.89	
104 Sweden	log(BR)	kg	cm	m	12–64	11–29	—	61	8	—	—	
105 Netherlands	CR	kg	cm	m	—	—	—	5	8	38	0.929	
106 Netherlands	CR	kg	cm	—	—	—	—	5	8	38	0.924	
107 UK	CR	t	cm	—	—	—	—	9	6	1	—	
108 UK	CR	t	cm	—	—	—	—	9	6	1	—	
109 France	ln(FL)	kg	cm	—	—	—	—	42	14	23	0.95	
110 Italy	FL	kg	cm	—	—	—	—	11	8	8	—	
111 Italy	FL	kg	cm	m	—	—	—	11	8	—	0.961	
112 Netherlands	FL	kg	cm	m	—	—	—	5	8	38	0.923	
113 Netherlands	FL	kg	cm	—	—	—	—	5	8	38	0.906	
114 Spain	FL	kg	cm	—	4–34.5	6.1–18.4	—	67	14	7	0.89	
115 France	ln(RC)	kg	cm	—	—	—	—	42	14	16	0.99	
116 France	ln(RF)	kg	cm	—	—	—	—	42	14	16	0.94	
117 France	ln(RS)	kg	cm	—	—	—	—	42	14	16	0.95	
118 France	RT	kg	cm	—	—	—	—	43	8	16	0.99	
119 France	log(RT)	kg	cm	—	3–20	—	—	21	8	16	0.99	
120 Germany	log(RT)	kg	cm	—	12–47	—	—	21	8	8	0.98	
121 Sweden	log(RT)	kg	cm	m	12–64	11–29	—	61	8	—	—	
122 France	ln(SB)	kg	cm	—	—	—	—	42	14	23	0.99	
123 Sweden	log(SB)	kg	cm	m	12–64	11–29	—	61	8	—	—	
124 Czech republic	ST	kg	cm	—	5.7–62.1	9.2–33.9	—	15	2	20	0.954	
125 Italy	ST	kg	cm	m	—	—	—	11	8	8	0.988	
126 Italy	ST	kg	cm	m	—	—	—	11	8	9	0.995	
127 Italy	ST	kg	cm	m	—	—	—	11	8	10	0.99	
128 Italy	ST	kg	cm	m	—	—	—	11	8	—	0.996	
129 Netherlands	ST	kg	cm	m	—	—	—	5	8	38	0.996	
130 Netherlands	ST	kg	cm	—	—	—	—	5	8	38	0.979	
131 Spain	ST	kg	cm	—	4–34.5	6.1–18.4	—	67	14	7	0.99	
132 France	ln(SW)	kg	cm	—	—	—	—	42	14	23	0.99	
133 Sweden	log(SW)	kg	cm	m	12–64	11–29	—	61	8	—	—	
<i>Fraxinus excelsior</i> (European ash)												
134 UK	ln(ABW)	kg	cm	—	2.9–33	—	—	10	14	11	15	0.994
135 UK	ln(ABW)	kg	cm	—	3–18	—	—	10	14	11	15	0.985
<i>Larix sibirica</i> (Siberian larch)												
136 Iceland	AB	kg	cm	m	3.3–31.6	3–20	—	71	8	44	0.992	
137 Iceland	ST	kg	cm	m	3.3–31.6	3–20	—	71	8	44	0.984	
<i>Larix</i> spp.												
138 UK	CR	t	cm	—	—	—	—	9	6	2	—	—
<i>Picea abies</i> (Norway spruce, Kuusi, Gran, Fichte, Rødgran, Epicéa)												
139 Belgium	AB	g	cm	cm	2.6–10	1.3–4.5	—	41	5	12	23	0.982
140 Czech republic	AB	kg	cm	m	1–11	2–9	—	18	7	13	55	—
141 Czech republic	AB	kg	cm	—	11–47	14–33	—	16	2	17	0.967	
142 Czech republic	AB	kg	cm	m	11–47	14–33	—	16	2	17	0.971	
143 Denmark	AB	kg	cm	m	10–17	11–13	—	59	7	14	5	—
144 Denmark	AB	kg	cm	m	12–20	11–14	—	59	7	15	10	—
145 Finland	AB	kg	cm	—	—	—	—	8	8	—	—	
146 Finland	AB	kg	cm	m	—	—	—	8	8	—	—	
147 Germany	AB	kg	cm	—	17–39	—	—	26	12	19	0.995	
148 Germany	AB	kg	cm	—	10–27.2	—	—	64	12	8	—	
149 Germany	AB	kg	cm	—	17–38	—	—	64	12	9	—	
150 Germany	AB	kg	cm	—	23–31	—	—	64	12	5	—	
151 Iceland	AB	kg	cm	m	2.7–27.9	2.7–12	—	71	8	16	0.981	
152 Norway	AB	gr	cm	—	5–15	—	—	6	7	35	0.993	
153 Norway	AB	gr	cm	—	2–5	—	—	6	7	35	—	
154 Sweden	log(AB)	kg	cm	m	15–38	18–28	—	61	8	—	—	
155 Belgium	log(ABW)	g	cm	—	16.–32.3	—	—	24	12	6	0.982	

Equation	a	b	Parameters		
			c	d	e
100	a+b·ln(D)	-6.2524	3.328	-	-
101	a·D <sup>b</sup> ·H <sup>c</sup>	0.0114	3.682	-1.031	-
102	a·D <sup>b</sup>	0.0020	3.265	-	-
103	a·D <sup>b</sup>	0.0317	2.3931	-	-
104	a+log[H·(D <sup>2</sup> )·b]	-3.2114	1.2481	-	-
105	a·D <sup>b</sup> ·H <sup>c</sup>	0.0183	3.614	-1.078	-
106	a·D <sup>b</sup>	0.0031	3.161	-	-
107	a·D <sup>2</sup>	2.595·10 <sup>-4</sup>	-	-	-
108	a+b·D <sup>c</sup>	0.00686	1.92·10 <sup>-5</sup>	2.4658	-
109	a+b·ln(D)	-4.8599	2.1935	-	-
110	a·D <sup>b</sup>	0.00295	2.43854	-	-
111	a·D <sup>b</sup> ·H <sup>c</sup>	0.02408	3.04567	-1.51571	-
112	a·D <sup>b</sup> ·H <sup>c</sup>	0.0167	2.951	-1.101	-
113	a+b·D <sup>c</sup>	0.375	0.0024	2.517	-
114	a·D <sup>b</sup>	0.0145	1.9531	-	-
115	a+b·ln(D)	-4.1302	2.6099	-	-
116	a+b·ln(D)	-5.7948	2.1609	-	-
117	a+b·ln(D)	-5.4415	2.082	-	-
118	a+b·ln(D)	-3.8219	2.5382	-	-
119	a+b·log(D)	-1.66	2.54	-	-
120	a+b·log(D)	-2	2.7	-	-
121	a+log[H·(D <sup>2</sup> )·b]	-2.8434	1.104	-	-
122	a+b·ln(D)	-3.0741	2.0543	-	-
123	a+log[H·(D <sup>2</sup> )·b]	-2.4279	0.8636	-	-
124	a·D <sup>b</sup>	0.494	2.07	-	-
125	a·D <sup>b</sup> ·H <sup>c</sup>	0.00519	1.49634	2.10419	-
126	a·D <sup>b</sup> ·H <sup>c</sup>	0.03638	2.15436	0.6587	-
127	a·D <sup>b</sup> ·H <sup>c</sup>	0.00269	2.02481	1.65219	-
128	a·D <sup>b</sup> ·H <sup>c</sup>	0.00519	1.87511	1.27233	-
129	a·D <sup>b</sup> ·H <sup>c</sup>	0.0109	1.951	1.262	-
130	a·D <sup>b</sup>	0.0762	2.523	-	-
131	a·D <sup>b</sup>	0.0894	2.4679	-	-
132	a+b·ln(D)	-2.0445	2.3912	-	-
133	a+log[H·(D <sup>2</sup> )·b]	-1.6219	0.9813	-	-
134	a+b·ln(D)	-2.4598	2.4882	-	-
135	a+b·ln(D)	-2.4718	2.5466	-	-
136	a·D <sup>b</sup> ·H <sup>c</sup>	0.1081	1.53	0.9482	-
137	a·D <sup>b</sup> ·H <sup>c</sup>	0.0444	1.4793	1.2397	-
138	a+b·D <sup>c</sup>	0.00564	3.041·10 <sup>-5</sup>	2.1058	-
139	a+b·H+c·D <sup>2</sup>	-520.7	-2.8	154.1	-
140	a·(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.4274	0.8674	1.0099	-0.2028
141	a·D <sup>b</sup>	0.57669	1.964	-	-
142	a·[(D <sup>2</sup> )·H] <sup>b</sup>	0.11975	0.81336	-	-
143	a·D <sup>2</sup> ·H	0.02155	-	-	-
144	a·D <sup>2</sup> ·H	0.01815	-	-	-
145	a+b·D+c·D <sup>2</sup>	19.018	-4.806	0.565	-
146	a+b·D <sup>2</sup> +c·(D <sup>2</sup> )·H	0.257	0.187	0.010	-
147	a+b·D+c·D <sup>2</sup>	-43.13	2.25	0.452	-
148	a+b·D+c·D <sup>2</sup>	-60.55702	5.46558	0.27567	-
149	a+b·D+c·D <sup>2</sup>	-283.17413	26.32334	-0.12856	-
150	a+b·D+c·D <sup>2</sup>	-142.60881	13.63896	0.12593	-
151	a·D <sup>b</sup> ·H <sup>c</sup>	0.2465	2.12	-0.167	-
152	a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	200.3691	99.3609	25	-
153	a·D <sup>2</sup>	200.3691	-	-	-
154	a+log[H·(D <sup>2</sup> )·b]	-1.2908	0.891	-	-
155	a+b·log(D)	1.81298	2.51353	-	-

## App. A

			Unit of Biom.	<i>D</i>	<i>H</i>	Range of <i>D</i> (cm)	Range of <i>H</i> (m)	Ref.	Cont.	Comm.	<i>n</i>	<i>r</i> <sup>2</sup>
156	Austria	BR	kg	mm	—	3.2–20.7	—	60	12	12	12	0.731
154	Austria	BR	kg	mm	—	2.5–17.9	—	60	12	12	12	0.95
158	Czech republic	BR	kg	cm	m	1–11	2–9	18	7	13	55	—
159	Czech republic	BR	kg	cm	—	11–47	14–33	16	2	25	25	0.892
160	Czech republic	BR	kg	cm	m	11–47	14–33	16	2	25	25	0.89
161	Denmark	ln(BR)	kg	cm	m	14–26	14–18	34	7	16	20	0.83
162	Europe	ln(BR)	kg	cm	—	1.8–67.6	2.1–42.8	74	8		429	0.871
163	Finland	BR	kg	cm	—	—	—	8	8	17	—	—
164	Finland	BR	kg	cm	m	—	—	8	8	17	—	—
165	Finland	BR	kg	cm	—	—	—	8	8	18	—	—
166	Finland	BR	kg	cm	m	—	—	8	8	18	—	—
167	Finland	BR	kg	cm	—	—	—	8	8	19	—	—
168	Finland	BR	kg	cm	m	—	—	8	8	19	—	—
169	Germany	BR	kg	cm	—	17–39	—	27	12	19	—	—
170	Germany	BR	kg	m	m	0–50	5–30	54	8	32	32	0.961
171	Germany	BR	kg	m	m	0–10	5–10	54	8	20	10	0.94
172	Germany	BR	kg	m	m	20–30	18–21	54	8	20	7	0.79
173	Germany	BR	kg	m	m	30–50	24–30	54	8	20	15	0.903
174	Germany	BR	kg	cm	—	26–60	30.–36.27	69	10	7	7	0.959
175	Germany	BR	kg	cm	—	27–55.8	22.–29.80	69	10	7	7	0.972
176	Germany	BR	kg	cm	—	23–52	32.–38.9	33	10	5	5	0.921
177	Germany	BR	kg	cm	—	26–60	30–36.26	69	10	20	7	0.892
178	Germany	BR	kg	cm	—	26–55.7	22–29.79	69	10	20	7	0.976
179	Germany	BR	kg	cm	—	23–52	32–38.8	33	10	5	5	0.886
180	Iceland	BR	kg	cm	m	2.7–27.9	2.7–12	71	8	4	16	0.944
181	Norway	BR	gr	cm	—	5–15	—	6	7	35	35	0.978
182	Norway	BR	gr	cm	—	2–5	—	6	7	35	—	—
183	Sweden	BR	kg	mm	—	4.9–29.8	4.1–23.4	37	4	32	32	0.955
184	Sweden	ln(BR)	g	cm	—	1.1–9.9	18–83	19	8	43	43	0.77
185	Sweden	log(BR)	kg	cm	m	15–38	18–28	61	8	—	—	—
186	Iceland	CR	kg	cm	m	2.7–27.9	2.7–12	71	8	16	16	0.959
187	Sweden	ln(CR)	kg	cm	—	0.3–63.4	1.3–35.6	49	8	1501	1501	0.933
188	Finland	CR	kg	cm	—	—	—	8	8	—	—	—
189	Finland	CR	kg	cm	m	—	—	8	8	—	—	—
190	Finland	CR	kg	mm	—	—	—	29	8	21	—	0.881
191	Finland	CR	kg	mm	dm	—	—	29	8	22	—	0.903
192	Finland	CR	kg	cm	—	—	—	28	8	—	—	0.912
193	Finland	CR	kg	mm	—	—	—	29	8	22	—	0.892
194	Finland	CR	kg	mm	dm	—	—	29	8	21	—	0.893
195	Germany	CR	kg	cm	—	26–60	—	69	12	7	7	0.964
196	Germany	CR	kg	cm	—	24–55.5	—	69	12	7	7	0.972
197	Germany	CR	kg	cm	—	10.–27.2	—	64	12	8	—	—
198	Germany	CR	kg	cm	—	17–38	—	64	12	9	—	—
199	Germany	CR	kg	cm	—	23–31	—	64	12	5	—	—
200	Germany	CR	kg	cm	—	24–52	32.–38.6	20	10	5	5	0.881
201	Sweden	ln(CR)	kg	cm	—	0–50	0–	50	8	544	544	0.945
202	Sweden	ln(CR)	kg	cm	m	0–50	0–	50	8	544	544	0.949
203	Belgium	log(DB)	g	cm	—	16.–32.3	—	24	12	6	6	0.97
204	Czech republic	DB	kg	cm	—	11–47	14–33	15	2	26	26	0.431
205	Czech republic	DB	kg	cm	m	11–47	14–33	15	2	26	26	0.41
206	Denmark	ln(DB)	kg	cm	m	14–26	14–18	34	7	16	20	0.58
207	Europe	ln(DB)	kg	cm	—	3.5–52.8	4.2–33.4	74	8		207	0.794
208	Finland	DB	kg	mm	—	—	—	28	8	—	—	0.266
209	Germany	DB	kg	cm	—	26–60	31–36	69	10	7	7	0.819
210	Norway	DB	gr	cm	—	5–15	—	6	7	35	35	0.798
211	Norway	DB	gr	cm	—	2–5	—	6	7	35	35	—
212	Sweden	ln(DB)	g	cm	dm	1.1–9.9	1.8–8.3	18	8	42	42	0.37
213	Sweden	ln(DB)	kg	cm	—	0–50	0–	50	8	525	525	0.714
214	Sweden	ln(DB)	kg	cm	m	0–50	0–	50	8	525	525	0.729
215	Sweden	ln(DB)	kg	cm	—	0.3–63.4	1.3–35.6	49	8	525	525	0.714

Equation	a	b	Parameters		
			c	d	e
156	a·exp(b·D)	5.3727	0.00876	—	—
157	a·exp(b·D)	1.325	0.0135	—	—
158	a·(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.1895	1.4075	0.8841	-0.9098
159	a·D <sup>b</sup>	0.0042229	2.7044	—	—
160	a·[(D <sup>2</sup> )·H] <sup>b</sup>	4.5394·10 <sup>-4</sup>	1.1262	—	—
161	a+b·ln(D <sup>2</sup> ·H)	-5.88	1.02	—	—
162	a+b·ln(D)	-3.96201	2.2552	—	—
163	a+b·D+c·D <sup>2</sup>	2.479	-0.552	0.066	—
164	a+b·D <sup>2</sup> +c·D·H	-0.129	0.076	-0.041	—
165	a+b·D+c·D <sup>2</sup>	3.640	-0.476	0.063	—
166	a+b·D+c·D <sup>2</sup> +d·D·H	3.225	-0.295	0.091	-0.045
167	a+b·D+c·D <sup>2</sup>	3.278	-0.728	0.101	—
168	a+b·D <sup>2</sup> +c·D·H	0.115	0.130	-0.074	—
169	a+b·D+c·D <sup>2</sup>	17.6	-2.87	0.141	—
170	a·(H·D <sup>2</sup> ) <sup>b</sup>	11.74	1.2	—	—
171	a·(H·D <sup>2</sup> ) <sup>b</sup>	33.53	0.916	—	—
172	a·(H·D <sup>2</sup> ) <sup>b</sup>	10.81	1.05	—	—
173	a·(H·D <sup>2</sup> ) <sup>b</sup>	3.98	1.306	—	—
174	a·D <sup>b</sup>	0.000135	3.453183	—	—
175	a·D <sup>b</sup>	1.96·10 <sup>-5</sup>	4.0576	—	—
176	a·D <sup>b</sup>	0.000999	2.833374	—	—
177	a·D <sup>b</sup>	0.0021994	2.4392	—	—
178	a·D <sup>b</sup>	0.0004947	2.9487	—	—
179	a·D <sup>b</sup>	0.00684	2.0603	—	—
180	a·D <sup>b</sup> ·H <sup>c</sup>	0.0653	2.9955	-1.3501	—
181	a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	37.7513	21.115	25	—
182	a·D <sup>2</sup>	37.7513	—	—	—
183	a·(1-exp(-b·D)) <sup>c</sup>	36.2826	0.0080	2.1576	—
184	a+b·ln(D) <sup>2</sup>	4.8678	0.8216	—	—
185	a+log[H·(D <sup>2</sup> )]·b	-3.628	1.2374	—	—
186	a·D <sup>b</sup> ·H <sup>c</sup>	0.2425	2.7517	-1.3456	—
187	a+b·[D/(D+13)]	-1.3858	8.6040	—	—
188	a+b·D+c·D <sup>2</sup>	3.416	-0.593	0.140	—
189	a+b·D+c·D <sup>2</sup> +d·H+e·(D·H)	6.721	0.358	0.119	-1.4773 0.030
190	a+b·D+c·D <sup>3</sup>	-3.71	0.10229	3.3·10 <sup>-6</sup>	—
191	a+b·D <sup>2</sup> +c·D <sup>3</sup> +d·(D <sup>3</sup> /H)	0.4866	3.5026·10 <sup>-4</sup>	1.35·10 <sup>-6</sup>	4.2424·10 <sup>-4</sup>
192	a+b·D <sup>2</sup>	3.2	0.1049	—	—
193	a+b·D+c·D <sup>3</sup>	-4.34	0.11571	3.34·10 <sup>-6</sup>	—
194	a+b·D <sup>2</sup> +c·D <sup>3</sup> +d·(D <sup>3</sup> /H)	0.4112	2.6724·10 <sup>-4</sup>	1.41·10 <sup>-6</sup>	4.3562·10 <sup>-4</sup>
195	a·D <sup>b</sup>	0.0013	3.1784	—	—
196	a·D <sup>b</sup>	0.0009	3.2112	—	—
197	a+b·D+c·D <sup>2</sup>	-0.55542	-0.39541	0.09537	—
198	a+b·D+c·D <sup>2</sup>	-70.51964	6.11247	-0.04391	—
199	a+b·D+c·D <sup>2</sup>	-12.29435	1.19256	0.0462	—
200	a·D <sup>b</sup>	0.1068	1.8137	—	—
201	a+b·[D/(D+13)]	-1.2804	8.5242	—	—
202	a+b·[D/(D+13)]+c·H+d·ln(H)	-1.2063	10.9708	-0.0124	-0.4923
203	a+b·log(D)	1.82795	1.49367	—	—
204	a·D <sup>b</sup>	0.055571	1.5317	—	—
205	a·[(D <sup>2</sup> )·H] <sup>b</sup>	0.021705	0.60715	—	—
206	a+b·ln(D <sup>2</sup> ·H)	-7.75	1.08	—	—
207	a+b·ln(D)	-3.22406	1.67320	—	—
208	a+b·D+c·D <sup>3</sup>	-0.62	0.0134	3.9·10 <sup>-8</sup>	—
209	a·D <sup>b</sup>	4.25·10 <sup>-7</sup>	4.731	—	—
210	a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	9.921	-3.6193	25	—
211	a·D <sup>2</sup>	9.921	—	—	—
212	a+b·ln(D <sup>2</sup> ·H)	-1.4358	0.7494	—	—
213	a+b·[D/(D+18)]	-4.3308	9.9550	—	—
214	a+b·[D/(D+18)]+c·H+d·ln(H)	-4.6351	3.6518	0.0493	1.0129
215	a+b·[D/(D+18)]	-4.6654	9.9550	—	—

## App. A

			Unit of Biom.	<i>D</i>	<i>H</i>	Range of <i>D</i> (cm)	Range of <i>H</i> (m)	Ref.	Cont.	Comm.	<i>n</i>	<i>r</i> <sup>2</sup>
216	Austria	FL	kg	mm	—	3.2–20.7	—	60	12	12	12	0.898
217	Austria	FL	kg	mm	—	2.5–17.9	—	60	12	12	12	0.89
218	Czech republic	FL	kg	cm	m	1–11	2–9	18	7	13	55	—
219	Czech republic	FL	kg	cm	—	11–47	14–33	16	2	25	25	0.941
220	Czech republic	FL	kg	cm	m	11–47	14–33	16	2	25	25	0.93
221	Denmark	ln(FL)	kg	cm	m	14–26	14–18	34	7	23	20	0.74
222	Europe	ln(FL)	kg	cm	—	1.8–67.6	2.1–42.8	74	8	24	551	0.847
223	Finland	FL	kg	cm	—	—	—	8	8	24	—	—
224	Finland	FL	kg	cm	m	—	—	8	8	24	—	—
225	Finland	ln(FL)	kg	mm	—	—	—	29	8	—	—	0.715
226	Finland	FL	kg	cm	m	—	—	46	8	—	—	—
227	Germany	log(FL)	kg	cm	—	18–62	—	70	12	28	28	0.911
228	Germany	FL	kg	cm	—	10–27.2	—	64	12	8	8	—
229	Germany	FL	kg	cm	—	17–38	—	64	12	9	9	—
230	Germany	FL	kg	m	m	0–10	5–10	54	8	10	10	0.961
231	Germany	FL	kg	m	m	20–30	18–21	54	8	7	7	0.869
232	Germany	FL	kg	m	m	30–50	24–30	54	8	15	15	0.906
233	Germany	FL	kg	cm	—	14–27	20–24	45	8	8	8	0.95
234	Germany	FL	kg	cm	—	26–60	30.–36.25	69	10	7	7	0.857
235	Germany	FL	kg	cm	—	25–55.6	22.–29.78	69	10	7	7	0.964
236	Germany	FL	kg	cm	—	24–52	32.–38.7	33	10	5	5	0.854
237	Norway	FL	gr	cm	—	5–15	—	6	7	35	35	0.978
238	Norway	FL	gr	cm	—	2–5	—	6	7	35	—	—
239	Sweden	ln(FL)	kg	cm	—	0–50	0–	50	8	544	544	0.899
240	Sweden	ln(FL)	kg	cm	m	0–50	0–	50	8	544	544	0.901
241	Sweden	ln(FL)	kg	cm	—	0.3–63.4	1.3–35.6	49	8	544	544	0.899
242	Sweden	FL	kg	mm	—	4.9–29.8	4.1–23.4	37	4	32	32	0.962
243	Sweden	ln(FL)	g	cm	—	1.1–9.9	18–83	19	8	43	43	0.87
244	Germany	RC	kg	m	m	0–50	5–30	54	8	25	25	0.872
245	Sweden	ln(RC)	kg	cm	—	0–50	0–	50	8	25	281	0.941
246	Germany	RS	kg	m	m	0–50	5–30	54	8	25	25	0.872
247	Sweden	ln(RS)	kg	cm	—	0–50	0–	50	8	26	329	0.925
248	Europe	ln(RT)	kg	cm	—	—	—	74	8	—	—	0.956
249	Germany	RT	kg	cm	—	15–23	—	22	12	15	15	0.63
250	Germany	RT	kg	cm	—	16–32.5	—	44	12	15	15	0.711
251	Germany	log(RT)	kg	cm	—	5–25	—	21	8	15	15	0.79
252	Germany	RT	kg	cm	—	14–27	20–24	45	8	5	5	0.96
253	Sweden	log(RT)	kg	cm	m	15–38	18–28	61	8	—	—	—
254	UK	RT	kg	cm	—	—	—	9	6	—	—	—
255	Czech republic	SB	kg	cm	—	11–47	14–33	16	2	18	18	0.927
256	Czech republic	SB	kg	cm	m	11–47	14–33	16	2	18	18	0.935
257	Denmark	ln(SB)	kg	cm	m	14–26	14–18	34	7	16	20	0.84
258	Germany	SB	kg	cm	—	10–27.2	—	64	12	8	8	—
259	Germany	SB	kg	cm	—	26–60	30.–36.28	69	10	7	7	0.965
260	Germany	SB	kg	cm	—	28–55.9	22.–29.81	69	10	7	7	0.862
261	Germany	SB	kg	cm	—	23–52	32.–38.10	33	10	5	5	0.904
262	Norway	SB	gr	cm	—	5–15	—	6	7	35	35	0.981
263	Norway	SB	gr	cm	—	2–5	—	6	7	35	—	—
264	Sweden	ln(SB)	kg	cm	—	0–50	0–	50	8	505	505	0.966
265	Sweden	ln(SB)	kg	cm	m	0–50	0–	50	8	505	505	0.968
266	Sweden	ln(SB)	kg	cm	—	0.3–63.4	1.3–35.6	49	8	505	505	0.966
267	Sweden	ln(SB)	g	cm	dm	1.1–9.9	1.8–8.3	19	8	43	43	0.97
268	Sweden	log(SB)	kg	cm	m	15–38	18–28	61	8	—	—	—
269	Sweden	ln(SR)	kg	cm	—	0–50	0–	50	8	316	316	0.97
270	Belgium	log(ST)	g	cm	—	16–32.3	—	24	12	6	6	0.986
271	Czech republic	ST	kg	cm	m	1–11	2–9	18	7	13	55	—
272	Europe	ln(ST)	kg	cm	—	3.5–52.8	4.2–33.4	74	8	235	235	0.986
273	Germany	ST	kg	cm	—	26–60	—	69	12	7	7	0.968
274	Germany	ST	kg	cm	—	24–55.5	—	69	12	7	7	0.962
275	Germany	ST	kg	cm	—	17–38	—	64	12	9	9	—

Equation	a	b	Parameters		
			c	d	e
216	a+b·D+c·D <sup>2</sup>	-1.9745	0.039	0.00382	-
217	a+b·D+c·D <sup>2</sup>	-0.7095	0.0011	0.00142	-
218	a·(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.2139	0.6896	1.2814	-0.6697
219	a·D <sup>b</sup>	0.030997	2.0019	-	-
220	a·[(D <sup>2</sup> )·H] <sup>b</sup>	0.0070936	0.81716	-	-
221	a+b·ln(D <sup>2</sup> ·H)	-4.85	0.81	-	-
222	a+b·ln(D)	-3.19632	1.91620	-	-
223	a+b·D+c·D <sup>2</sup>	-0.647	0.106	0.040	-
224	a+b·D+c·D <sup>2</sup> +d·D·H	-0.607	0.095	0.039	0.001
225	a+b·ln(D)	-9.03	2.2204	-	-
226	a·D <sup>b</sup> ·H <sup>c</sup>	0.1022	2.5947	-0.8647	-
227	a+b·log(D)	-3.084	2.814	-	-
228	a+b·D+c·D <sup>2</sup>	-12.29769	1.14647	0.00179	-
229	a+b·D+c·D <sup>2</sup>	-12.55702	1.14647	0.00179	-
230	a·(H·D <sup>2</sup> ) <sup>b</sup>	35.05	0.847	-	-
231	a·(H·D <sup>2</sup> ) <sup>b</sup>	21.37	0.979	-	-
232	a·(H·D <sup>2</sup> ) <sup>b</sup>	9.26	1.179	-	-
233	a+b·D	-18.63	1.85	-	-
234	a·D <sup>b</sup>	0.0061379	2.3026	-	-
235	a·D <sup>b</sup>	0.0026146	2.6763	-	-
236	a·D <sup>b</sup>	0.00784	2.14426	-	-
237	a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	53.0637	10.4186	25	-
238	a·D <sup>2</sup>	53.0637	-	-	-
239	a+b·[D/(D+12)]	-1.9602	7.8171	-	-
240	a+b·[D/(D+12)]+c·ln(H)	-1.8551	9.7809	-0.4873	-
241	a+b·[D/(D+12)]	-2.0330	7.8171	-	-
242	a·(1-exp(-b·D)) <sup>c</sup>	348.6448	0.0025	2.6100	-
243	a+b·ln(D) <sup>2</sup>	5.5129	0.7519	-	-
244	a·(H·D <sup>2</sup> ) <sup>b</sup>	7.33	1.383	-	-
245	a+b·[D/(D+8)]	-6.3851	13.3703	-	-
246	a·(H·D <sup>2</sup> ) <sup>b</sup>	1.13	0.926	-	-
247	a+b·[D/(D+12)]	-2.5706	7.6283	-	-
248	a+b·ln(D)	-5.37891	2.92111	-	-
249	a·D <sup>b</sup>	0.02	2.36	-	-
250	a+b·D	-33.225	2.3915	-	-
251	a+b·log(D)	-1.7	2.36	-	-
252	a+b·D	-45.94	3.58	-	-
253	a+log[H·(D <sup>2</sup> )·b]	-2.0274	0.8946	-	-
254	a·D <sup>b</sup>	1.204·10 <sup>-5</sup>	2.4920	-	-
255	a·D <sup>b</sup>	0.032777	1.8902	-	-
256	a·[(D <sup>2</sup> )·H] <sup>b</sup>	0.0071913	0.7832	-	-
257	a+b·ln(D <sup>2</sup> ·H)	-5.51	0.88	-	-
258	a+b·D+c·D <sup>2</sup>	-6.55127	0.75517	0.02156	-
259	a·D <sup>b</sup>	0.23943	1.439	-	-
260	a·D <sup>b</sup>	0.2917	1.2919	-	-
261	a·D <sup>b</sup>	0.1557	1.5908	-	-
262	a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	23.8849	-3.9241	25	-
263	a·D <sup>2</sup>	23.8849	-	-	-
264	a+b·[D/(D+15)]	-3.3912	9.8364	-	-
265	a+b·[D/(D+15)]+c·H+d·ln(H)	-3.4020	8.3089	0.0147	0.2295
266	a+b·[D/(D+15)]	-3.4216	9.8364	-	-
267	a+b·ln(D)+c·H+d·ln[(D <sup>2</sup> )·H]	7.0429	4.946	0.0438	-1.5405
268	a+log[H·(D <sup>2</sup> )·b]	-1.8073	0.7426	-	-
269	a+b·[D/(D+14)]	-2.4447	10.5381	-	-
270	a+b·log(D)	1.85007	2.45530	-	-
271	a·(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.0468	1.2244	0.5465	0.6481
272	a+b·ln(D)	-2.50602	2.44277	-	-
273	a·D <sup>b</sup>	0.408	2.0136	-	-
274	a·D <sup>b</sup>	0.208	2.1531	-	-
275	a+b·D+c·D <sup>2</sup>	-212.04143	20.20032	-0.08466	-

## App. A

			Unit of Biom.	D	H	Range of D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	$r^2$
276	Germany	ST	kg	cm	—	23–31	—	64	12	5	—	
277	Germany	ST	ton	cm	—	17–39	21–31	26	7	19	0.989	
278	Germany	ST	kg	cm	—	24–52	32.–38.5	20	10	5	0.984	
279	Iceland	ST	kg	cm	m	2.7–27.9	2.7–12	71	8	16	0.989	
280	Sweden	ln(ST)	kg	cm	—	0–50	0–	50	8	546	0.988	
281	Sweden	ln(ST)	kg	cm	m	0–50	0–	50	8	546	0.994	
282	Sweden	ln(ST)	kg	cm	m	0.3–63.4	1.3–35.6	49	8	1503	0.994	
283	Sweden	ln(ST)	kg	cm	—	0.3–63.4	1.3–35.6	49	8	1503	0.986	
284	Sweden	ln(ST)	kg	cm	m	0.3–63.4	1.3–35.6	49	8	505	0.992	
285	Sweden	ln(ST)	kg	cm	—	0.3–63.4	1.3–35.6	49	8	505	0.982	
286	Sweden	ln(SU)	kg	cm	—	0–50	0–	50	8	328	0.958	
287	Czech republic	SW	kg	cm	—	11–47	14–33	16	2	18	0.969	
288	Czech republic	SW	kg	cm	—	11–47	14–33	16	2	18	0.976	
289	Denmark	ln(SW)	kg	cm	m	14–26	14–18	34	7	16	20	0.96
290	Germany	SW	kg	cm	—	26–60	30.–36.29	69	10	7	0.956	
291	Germany	SW	kg	cm	—	29–55.1	22.–29.82	69	10	7	0.963	
292	Germany	SW	kg	cm	—	23–51	32.–38.11	33	10	5	0.984	
293	Norway	SW	gr	cm	—	5–15	—	6	7	35	0.985	
294	Norway	SW	gr	cm	—	2–5	—	6	7	35	—	
295	Sweden	ln(SW)	kg	cm	—	0–50	0–	50	8	505	0.982	
296	Sweden	ln(SW)	kg	cm	m	0–50	0–	50	8	505	0.992	
297	Sweden	ln(SW)	g	cm	dm	1.1–9.9	1.8–8.3	19	8	43	0.99	
298	Sweden	log(SW)	kg	cm	m	15–38	18–28	61	8	—	—	
299	Germany	TB	kg	cm	—	26–60	30.–36.23	20	10	7	0.974	
300	Germany	TB	kg	cm	—	24–55.5	22.–29.77	20	10	7	0.975	
301	Germany	TB	kg	cm	—	24–52	32.–38.4	20	10	5	0.981	
<i>Picea engelmannii</i> (Engelmann spruce)												
302	Iceland	AB	kg	cm	m	1.4–12.7	1.7–12.7	71	8	14	0.927	
303	Iceland	ST	kg	cm	m	1.4–12.7	1.7–12.7	71	8	14	0.967	
<i>Picea rubens</i>												
304 –		log(RT)	kg	cm	—	—	—	68	8	15	0.972	
<i>Picea sitchensis</i> (Sitka spruce)												
305	UK	RT	kg	cm	—	—	—	9	6	—	—	
<i>Picea</i> spp.												
306	Iceland	AB	kg	cm	m	4.9–28.6	4.8–15.4	71	8	56	0.965	
307	Iceland	CR	kg	cm	m	4.9–28.6	4.8–15.4	71	8	56	0.905	
308	UK	CR	kg	cm	—	—	—	9	6	1	—	
309	UK	CR	kg	cm	—	—	—	9	6	2	—	
310	Iceland	ST	kg	cm	m	4.9–28.6	4.8–15.4	71	8	56	0.981	
<i>Pinus banksiana</i>												
311 –		log(RT)	kg	cm	—	—	—	68	8	40	0.917	
<i>Pinus contorta</i> (Lodgepole pine)												
312	Iceland	AB	kg	cm	m	4.2–26.3	2.8–12.8	71	8	48	0.984	
313	UK	CR	kg	cm	—	—	—	9	6	2	—	
314 –		log(RT)	kg	cm	m	—	—	68	8	72	0.949	
315 –		log(RT)	kg	cm	m	—	—	68	8	221	0.9	
316	UK	RT	kg	cm	—	—	—	9	6	—	—	
317	Iceland	ST	kg	cm	m	4.2–26.3	2.8–12.8	71	8	48	0.992	
<i>Pinus nigra</i> var <i>maritima</i> (Black pine, Corsican pine)												
318	UK	CR	kg	cm	—	—	—	9	6	2	—	
319	UK	RT	kg	cm	—	—	—	9	6	—	—	
<i>Pinus pinaster</i> (Maritime pine)												
320	Italy	ln(AB)	kg	cm	—	1.5–16	—	2	14	6	8	0.99
<i>Pinus radiata</i> (Radiata pine)												
321	Italy	ln(AB)	kg	cm	—	4–20	—	53	14	6	17	0.99
322 –		log(RT)	kg	cm	—	—	—	68	8	8	0.944	
323 –		log(RT)	kg	cm	m	—	—	68	8	8	0.943	
<i>Pinus sylvestris</i> (Scots pine, Mänty, Tall)												
324	Czech republic	AB	kg	cm	m	2–6	2–5	17	7	29	—	
325	Czech republic	AB	kg	cm	m	2–16	4–11	17	7	50	—	

	Equation	a	b	Parameters		
				c	d	e
276	a+b·D+c·D <sup>2</sup>	-784.923	61.58581	-0.79535	-	-
277	a+b·D+c·D <sup>2</sup>	0.051	0.0038	0.000344	-	-
278	a·D <sup>b</sup>	0.5938	1.9423	-	-	-
279	a·D <sup>b</sup> ·H <sup>c</sup>	0.0712	1.637	0.7436	-	-
280	a+b·[D/(D+14)]	-2.0571	11.3341	-	-	-
281	a+b·[D/(D+14)]+c·H+d·ln(H)	-2.1702	7.4690	0.0289	0.6858	-
282	a+b·[D/(D+14)]+c·H+d·ln(H)	-2.2052	7.4361	0.0186	0.7595	-
283	a+b·[D/(D+14)]	-2.0148	11.1926	-	-	-
284	a+b·[D/(D+14)]+c·H+d·ln(H)	-2.3036	7.2309	0.0355	0.7030	-
285	a+b·[D/(D+14)]	-2.2727	11.4873	-	-	-
286	a+b·[D/(D+17)]	-3.3645	10.6686	-	-	-
287	a·D <sup>b</sup>	0.52917	1.9123	-	-	-
288	a·[(D <sup>2</sup> )·H] <sup>b</sup>	0.115	0.79159	-	-	-
289	a+b·ln(D <sup>2</sup> ·H)	-3.24	0.88	-	-	-
290	a·D <sup>b</sup>	0.31974	2.0595	-	-	-
291	a·D <sup>b</sup>	0.15739	2.2118	-	-	-
292	a·D <sup>b</sup>	0.5937753	1.9423097	-	-	-
293	a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	75.7482	75.3706	25	-	-
294	a·D <sup>2</sup>	75.7482	-	-	-	-
295	a+b·[D/(D+14)]	-2.2471	11.4873	-	-	-
296	a+b·[D/(D+14)]+c·H+d·ln(H)	-2.3032	7.2309	0.0355	0.7030	-
297	a+b·ln(D) <sup>2</sup> +c·ln[(D <sup>2</sup> )·H]	3.6562	0.4115	0.401	-	-
298	a+log[H·(D <sup>2</sup> )·b]	-1.2187	0.8494	-	-	-
299	a·D <sup>b</sup>	0.2543	2.1872	-	-	-
300	a·D <sup>b</sup>	0.1245	2.3585	-	-	-
301	a·D <sup>b</sup>	0.8007	1.9075	-	-	-
302	a·D <sup>b</sup> ·H <sup>c</sup>	0.9211	1.438	0.102	-	-
303	a·D <sup>b</sup> ·H <sup>c</sup>	0.2288	1.239	0.717	-	-
304	a·log(D)+b	2.1514	-1.2417	-	-	-
305	a·D <sup>b</sup>	1.115·10 <sup>-5</sup>	2.68358	-	-	-
306	a·D <sup>b</sup> ·H <sup>c</sup>	0.1334	1.8716	0.4386	-	-
307	a·D <sup>b</sup> ·H <sup>c</sup>	0.087	2.287	-0.2897	-	-
308	a·D <sup>b</sup>	5.2193·10 <sup>-4</sup>	1.459	-	-	-
309	a+b·D <sup>c</sup>	0.0060722	9.58·10 <sup>-6</sup>	2.5578	-	-
310	a·D <sup>b</sup> ·H <sup>c</sup>	0.0558	1.5953	0.9336	-	-
311	a·log(D)+b	2.16	-0.2089	-	-	-
312	a·D <sup>b</sup> ·H <sup>c</sup>	0.1429	1.8887	0.4332	-	-
313	a+b·D <sup>c</sup>	0.00435	1.321·10 <sup>-5</sup>	2.5138	-	-
314	a·log(H·D <sup>2</sup> )+b	1.022	-1.818	-	-	-
315	a·log(H·D <sup>2</sup> )+b	0.806	-1.062	-	-	-
316	a·D <sup>b</sup>	2.242·10 <sup>-5</sup>	2.42909375	-	-	-
317	a·D <sup>b</sup> ·H <sup>c</sup>	0.0669	1.5958	0.9096	-	-
318	a·D <sup>b</sup>	1.3997·10 <sup>-4</sup>	1.72105599	-	-	-
319	a·D <sup>b</sup>	1.537·10 <sup>-5</sup>	2.39136175	-	-	-
320	a+b·ln(D)	-1.457	1.8647	-	-	-
321	a+b·ln(D)	-2.359	2.2936	-	-	-
322	a·log(D)+b	2.4453	-0.9366	-	-	-
323	a·log(H·D <sup>2</sup> )+b	1.0519	-2.9005	-	-	-
324	a(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.0398	2.2993	0.4445	0.1232	-
325	a(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.0146	2.3868	-0.0618	0.8581	-

## App. A

		Biom.	Unit of D	H	Range of D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	$r^2$
326	Czech republic	AB	kg	cm	—	2–6	2–5	17	8	—	0.91
327	Czech republic	AB	kg	cm	—	2–6	2–5	17	8	—	0.91
328	Czech republic	AB	kg	cm	—	2–6	2–5	17	8	—	0.94
329	Czech republic	AB	kg	cm	m	2–6	2–5	17	8	—	0.47
330	Czech republic	AB	kg	cm	m	2–6	2–5	17	8	—	0.45
331	Czech republic	AB	kg	cm	m	2–6	2–5	17	8	—	0.45
332	Czech republic	AB	kg	cm	—	2–16	4–11	17	8	—	0.89
333	Czech republic	AB	kg	cm	—	2–16	4–11	17	8	—	0.67
334	Czech republic	AB	kg	cm	—	2–16	4–11	17	8	—	0.98
335	Czech republic	AB	kg	cm	m	2–16	4–11	17	8	—	0.52
336	Czech republic	AB	kg	cm	m	2–16	4–11	17	8	—	0.53
337	Czech republic	AB	kg	cm	m	2–16	4–11	17	8	—	0.82
338	Finland	AB	kg	cm	—	—	—	8	8	—	—
339	Finland	AB	kg	cm	—	—	—	8	8	—	—
340	Finland	ln(AB)	kg	cm	—	—	—	55	8	28	18
341	Finland	ln(AB)	kg	cm	—	—	—	55	8	29	30
342	Norway	AB	gr	cm	—	7–15	—	6	7	16	0.993
343	Norway	AB	gr	cm	—	2–7	—	6	7	16	—
344	Norway	AB	gr	cm	—	7–20	—	6	7	30	16
345	Norway	AB	gr	cm	—	2–7	—	6	7	30	16
346	Poland	ln(AB)	kg	cm	—	—	—	62	8	65	0.764
347	Poland	ln(AB)	kg	cm	—	—	—	62	8	110	0.89
348	Poland	ln(AB)	kg	cm	—	—	—	62	8	30	0.938
349	Poland	ln(AB)	kg	cm	—	—	—	62	8	15	0.975
350	UK	log(AB)	g	cm	—	—	—	48	8	10	0.996
351	Poland	ln(ABW)	kg	cm	—	—	—	62	8	65	0.775
352	Poland	ln(ABW)	kg	cm	—	—	—	62	8	110	0.894
353	Poland	ln(ABW)	kg	cm	—	—	—	62	8	30	0.938
354	Poland	ln(ABW)	kg	cm	—	—	—	62	8	15	0.978
355	Belgium	BR	kg	cm	—	—	—	75	13	—	—
356	Czech republic	BR	kg	cm	m	2–6	2–5	17	7	29	—
357	Czech republic	BR	kg	cm	m	2–16	4–11	17	7	50	—
358	Czech republic	BR	kg	cm	—	2–6	2–5	17	8	—	0.81
359	Czech republic	BR	kg	cm	—	2–6	2–5	17	8	—	0.83
360	Czech republic	BR	kg	cm	—	2–6	2–5	17	8	—	0.81
361	Czech republic	BR	kg	cm	—	2–16	4–11	17	8	—	0.77
362	Czech republic	BR	kg	cm	—	2–16	4–11	17	8	—	0.71
363	Czech republic	BR	kg	cm	—	2–16	4–11	17	8	—	0.89
364	Czech republic	BR	kg	cm	m	2–16	4–11	17	8	—	0.74
365	Czech republic	BR	kg	cm	m	2–16	4–11	17	8	—	0.67
366	Finland	BR	kg	cm	—	—	—	8	8	17	—
367	Finland	BR	kg	cm	m	—	—	8	8	17	—
368	Finland	BR	kg	cm	—	—	—	8	8	18	—
369	Finland	BR	kg	cm	m	—	—	8	8	18	—
370	Finland	BR	kg	cm	—	—	—	8	8	19	—
371	Finland	BR	kg	cm	m	—	—	8	8	19	—
372	Finland	ln(BR)	kg	cm	—	—	—	55	8	28	18
373	Finland	ln(BR)	kg	cm	—	—	—	55	8	29	30
374	Norway	BR	gr	cm	—	7–15	—	6	7	16	0.974
375	Norway	BR	gr	cm	—	2–7	—	6	7	16	—
376	Norway	BR	gr	cm	—	15–20	—	6	7	30	16
377	Norway	BR	gr	cm	—	2–15	—	6	7	30	16
378	Sweden	BR	kg	cm	m	2–40	2–	1	8	73	0.49
379	Sweden	ln(BR)	g	cm	dm	1.1–10	1.7–8.8	19	8	84	0.73
380	UK	log(BR)	g	cm	—	—	—	48	8	10	0.964
381	Norway	CO	gr	cm	—	7–15	—	6	7	16	0.688
382	Norway	CO	gr	cm	—	2–7	—	6	7	16	—
383	Norway	CO	gr	cm	—	2–13	—	6	7	30	16
384	Finland	CR	kg	cm	—	—	—	8	8	—	—
385	Finland	CR	kg	cm	m	—	—	8	8	—	—

Equation	a	b	Parameters		
			c	d	e
326 a+b·D	-3.1381	1.7295	-	-	-
327 a-exp(D·b)	0.2304	0.6536	-	-	-
328 a·D <sup>b</sup>	0.1599	2.2060	-	-	-
329 a+b·H	-2.2818	1.3799	-	-	-
330 a-exp(H·b)	0.332	0.51	-	-	-
331 a·H <sup>b</sup>	0.2146	1.8238	-	-	-
332 a+b·D	-25.2864	5.4433	-	-	-
333 a-exp(D·b)	1.8829	0.2445	-	-	-
334 a·D <sup>b</sup>	0.1182	2.3281	-	-	-
335 a+b·H	-53.8304	8.7802	-	-	-
336 a-exp(H·b)	0.591	0.3794	-	-	-
337 a·H <sup>b</sup>	0.0023	4.1398	-	-	-
338 a+b·D+c·D <sup>2</sup>	18.779	-4.328	0.506	-	-
339 a+b·D+c·D <sup>2</sup>	7.041	-1.279	0.201	-	-
340 a+b·ln(D)	-3.2807	2.6931	-	-	-
341 a+b·ln(D)	-2.3042	2.2608	-	-	-
342 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	209.69901	48.8075	49	-	-
343 a·D <sup>2</sup>	209.69901	-	-	-	-
344 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	200.87186	124.6808	49	-	-
345 a·D <sup>2</sup>	200.87186	-	-	-	-
346 a+b·ln(D)	-1.954	1.988	-	-	-
347 a+b·ln(D)	-2.202	2.112	-	-	-
348 a+b·ln(D)	-2.103	1.994	-	-	-
349 a+b·ln(D)	-2.001	1.943	-	-	-
350 a+b·log( $\pi$ ·D)	0.981	2.289	-	-	-
351 a+b·ln(D)	-1.979	1.959	-	-	-
352 a+b·ln(D)	-2.204	2.069	-	-	-
353 a+b·ln(D)	-2.087	1.939	-	-	-
354 a+b·ln(D)	-2.017	1.915	-	-	-
355 a·D <sup>b</sup>	0.0022	2.9122	-	-	-
356 a·(D+1) <sup>b+c·log(D)^9</sup> ·H <sup>d</sup>	0.1147	-0.585	3.1296	-0.4967	-
357 a·(D+1) <sup>b+c·log(D)^9</sup> ·H <sup>d</sup>	0.0228	-0.2728	1.8144	0.6324	-
358 a+b·D	-0.9086	0.4691	-	-	-
359 a-exp(D·b)	0.0493	0.6818	-	-	-
360 a·D <sup>b</sup>	0.0356	2.2530	-	-	-
361 a+b·D	-8.9086	1.5897	-	-	-
362 a-exp(D·b)	0.1862	0.2989	-	-	-
363 a·D <sup>b</sup>	0.0071	2.7743	-	-	-
364 a-exp(H·b)	0.0055	0.7173	-	-	-
365 a·H <sup>b</sup>	0.0001	4.6817	-	-	-
366 a+b·D+c·D <sup>2</sup>	2.842	-0.725	0.060	-	-
367 a+b·D+c·D <sup>2</sup> +d·D·H	3.129	-0.536	0.077	-0.036642	-
368 a+b·D+c·D <sup>2</sup>	2.171	1.599	0.060	-	-
369 a+b·D+c·D <sup>2</sup> +d·H	-0.398	-0.642	0.116	1.0275	-
370 a+b·D+c·D <sup>2</sup>	4.988	-1.104	0.087	-	-
371 a+b·D+c·D <sup>2</sup> +d·H+e·(D·H)	-2.589	-0.588	0.145	1.1308	-0.143
372 a+b·ln(D)	-3.1296	2.0089	-	-	-
373 a+b·ln(D)	-6.3284	3.0119	-	-	-
374 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	59.98243	-8.2237	49	-	-
375 a·D <sup>2</sup>	59.98243	-	-	-	-
376 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	32.9683	-8.27642	225	-	-
377 a·D <sup>2</sup>	32.9683	-	-	-	-
378 exp(a)·(D+1) <sup>b</sup> ·H <sup>c</sup>	-3.391	3.263	-1.202	-	-
379 a+b·ln(D)+c·ln(D <sup>2</sup> )+d·ln(H)	7.0826	1.2301	0.5838	-1.139	-
380 a+b·log( $\pi$ ·D)	-0.662	2.768	-	-	-
381 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	2.32445	-1.628	49	-	-
382 a·D <sup>2</sup>	2.32445	-	-	-	-
383 a·D <sup>2</sup>	0.27462	-	-	-	-
384 a+b·D+c·D <sup>2</sup>	8.033	-1.156	0.105	-	-
385 a+b·D+c·D <sup>2</sup> +d·H+e·(D·H)	-1.057	-0.506	0.176	1.3419	-0.173

## App. A

		Unit of Biom.	D	H	Range of D (cm)		H (m)		Ref.	Cont.	Comm.	n	$r^2$
386 Finland	ln(CR)	kg	mm	—	—	—	29	8	22	—	—	0.905	
387 Finland	ln(CR)	kg	mm	m	—	—	29	8	22	—	—	0.925	
388 Finland	CR	kg	cm	—	—	—	28	8	—	—	—	0.91	
389 Finland	ln(CR)	kg	mm	—	—	—	29	8	21	—	—	0.878	
390 Finland	ln(CR)	kg	mm	m	—	—	29	8	21	—	—	0.908	
391 Sweden	ln(CR)	kg	cm	—	0–45	0–	50	8	—	482	—	0.901	
392 Sweden	ln(CR)	kg	cm	m	0–45	0–	50	8	—	482	—	0.922	
393 UK	log(CR)	g	cm	—	—	—	48	8	—	10	—	0.975	
394 UK	CR	kg	cm	—	—	—	9	6	2	—	—	—	
395 Finland	log(DB)	kg	cm	m	2.4–9.7	3–7.8	56	8	—	—	—	0.742	
396 Finland	log(DB)	kg	cm	m	7–21.6	7.9–14.8	56	8	—	—	—	0.655	
397 Finland	log(DB)	kg	cm	m	5.6–23.6	8–18.2	56	8	—	—	—	0.89	
398 Finland	DB	kg	mm	—	—	—	29	8	—	—	—	0.228	
399 Norway	DB	gr	cm	—	2–7	—	6	7	30	16	—	—	
400 Norway	DB	gr	cm	—	7–20	—	6	7	30	16	0.925	—	
401 Norway	DB	gr	cm	—	11–15	—	6	7	31	16	0.989	—	
402 Norway	DB	gr	cm	—	2–11	—	6	7	—	16	—	—	
403 Sweden	ln(DB)	kg	cm	—	0–45	0–	50	8	—	467	0.741	—	
404 Sweden	ln(DB)	kg	cm	m	0–45	0–	50	8	—	467	0.748	—	
405 Sweden	ln(DB)	g	cm	dm	1.1–10	1.7–8.8	19	8	—	80	0.58	—	
406 UK	log(DB)	g	cm	—	—	—	48	8	—	10	0.512	—	
407 Belgium	FL	kg	cm	—	—	—	75	13	—	—	—	—	
408 Czech republic	FL	kg	cm	m	2–6	2–5	17	7	32	29	—	—	
409 Czech republic	FL	kg	cm	m	2–16	4–11	17	7	—	50	—	—	
410 Czech republic	FL	kg	cm	—	2–6	2–5	17	8	—	—	0.67	—	
411 Czech republic	FL	kg	cm	—	2–6	2–5	17	8	—	—	0.68	—	
412 Czech republic	FL	kg	cm	—	2–6	2–5	17	8	—	—	0.72	—	
413 Czech republic	FL	kg	cm	—	2–16	4–11	17	8	—	—	0.84	—	
414 Czech republic	FL	kg	cm	—	2–16	4–11	17	8	—	—	0.8	—	
415 Czech republic	FL	kg	cm	—	2–16	4–11	17	8	—	—	0.89	—	
416 Czech republic	FL	kg	cm	m	2–16	4–11	17	8	—	—	0.43	—	
417 Czech republic	FL	kg	cm	m	2–16	4–11	17	8	—	—	0.75	—	
418 Czech republic	FL	kg	cm	m	2–16	4–11	17	8	—	—	0.72	—	
419 Finland	FL	kg	cm	—	—	—	8	8	24	—	—	—	
420 Finland	FL	kg	cm	—	—	—	8	8	24	—	—	—	
421 Finland	ln(FL)	kg	cm	—	—	—	55	8	28	18	0.78	—	
422 Finland	ln(FL)	kg	cm	—	—	—	55	8	29	30	0.9	—	
423 Finland	ln(FL)	kg	mm	—	—	—	29	8	—	—	0.688	—	
424 Finland	FL	kg	cm	m	—	—	46	8	—	—	—	—	
425 Norway	FL	gr	cm	—	7–15	—	6	7	—	16	0.95	—	
426 Norway	FL	gr	cm	—	2–7	—	6	7	—	16	—	—	
427 Norway	FL	gr	cm	—	7–20	—	6	7	30	16	0.946	—	
428 Norway	FL	gr	cm	—	2–7	—	6	7	30	16	—	—	
429 Poland	ln(FL)	kg	cm	—	—	—	62	8	—	64	0.355	—	
430 Poland	ln(FL)	kg	cm	—	—	—	62	8	—	108	0.546	—	
431 Poland	ln(FL)	kg	cm	—	—	—	62	8	—	29	0.752	—	
432 Poland	FL	kg	cm	cm	0.2–9.2	1.3–5.6	3	8	—	—	0.75	—	
433 Poland	ln(FL)	kg	cm	—	—	—	62	8	—	15	0.71	—	
434 Sweden	ln(FL)	kg	cm	—	0–45	0–	50	8	—	482	0.841	—	
435 Sweden	ln(FL)	kg	cm	m	0–45	0–	50	8	—	482	0.865	—	
436 Sweden	FL	kg	cm	m	2–40	2–	1	8	—	73	0.908	—	
437 Sweden	ln(FL)	g	cm	dm	1.1–10	1.7–8.8	19	8	—	84	0.75	—	
438 UK	log(FL)	g	cm	—	—	—	48	8	—	10	0.962	—	
439 Belgium	FL(1)	kg	cm	—	—	—	75	13	33	—	—	—	
440 Belgium	FL(2)	kg	cm	—	—	—	75	13	34	—	—	—	
441 Belgium	RC	kg	cm	—	—	—	75	13	—	—	—	—	
442 Europe	RC	kg	cm	—	4.1–45	—	77	1	35	—	0.97	—	
443 Poland	ln(RC)	kg	cm	—	—	—	62	8	—	61	0.589	—	
444 Poland	ln(RC)	kg	cm	—	—	—	62	8	—	106	0.803	—	
445 Poland	ln(RC)	kg	cm	—	—	—	62	8	—	30	0.891	—	

Equation	a	b	Parameters		
			c	d	e
386 a+b·ln(D)	-8.8027	2.2475	—	—	—
387 a+b·D+c·ln(D)+d·H	-9.7486	0.0016023	2.5600	-0.0063173	—
388 a+b·D <sup>2</sup>	2.7	0.0799	—	—	—
389 a+b·ln(D)	-9.3954	2.3268	—	—	—
390 a+b·ln(D)+c·ln(H)+d·D/H <sup>2</sup>	-5.2678	3.4914	-1.9498	-47.454	—
391 a+b·[D/(D+10)]	-2.8604	9.1015	—	—	—
392 a+b·[D/(D+10)]+c·ln(H)	-2.5413	13.3955	-1.1955	—	—
393 a+b·log(π·D)	-0.383	2.723	—	—	—
394 a+b·D <sup>c</sup>	0.00435122	1.321·10 <sup>-5</sup>	2.51380074	—	—
395 a+b·log(D <sup>2</sup> ·H)	-2.130	0.6797	—	—	—
396 a+b·log(D <sup>2</sup> ·H)	-2.702	0.8486	—	—	—
397 a+b·log(D <sup>2</sup> ·H)	-2.735	0.8977	—	—	—
398 a+b·D	-0.84	0.0194	—	—	—
399 a·D <sup>2</sup>	10.3135	—	—	—	—
400 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	10.3135	15.8006	49	—	—
401 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	4.21541	19.4571	121	—	—
402 a·D <sup>2</sup>	4.21541	—	—	—	—
403 a+b·[D/(D+10)]	-5.3338	9.5938	—	—	—
404 a+b·[D/(D+10)]+c·H+d·ln(H)	-5.8926	7.1270	-0.0465	1.1060	—
405 a+b·ln(D)+c·H	1.1059	0.7017	0.0494	—	—
406 a+b·log(π·D)	0.959	1.569	—	—	—
407 a·D <sup>b</sup>	0.00445	2.2371	—	—	—
408 a·(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.0009	6.7765	-2.1552	-1.3905	—
409 a·(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.0012	2.6479	-0.0549	0.6776	—
410 a+b·D	-0.564	0.3324	—	—	—
411 a-exp(D·b)	0.0425	0.6652	—	—	—
412 a·D <sup>b</sup>	0.0279	2.2877	—	—	—
413 a+b·D	-2.9979	0.6293	—	—	—
414 a-exp(D·b)	0.1044	0.3192	—	—	—
415 a·D <sup>b</sup>	0.009	2.4794	—	—	—
416 a+b·H	-5.6616	0.9373	—	—	—
417 a-exp(H·b)	0.0066	0.6513	—	—	—
418 a·H <sup>b</sup>	0.0002	4.3542	—	—	—
419 a·D+b·D <sup>2</sup>	0.023	0.015	—	—	—
420 a+b·D+c·D <sup>2</sup>	-0.105	0.365	0.010	—	—
421 a+b·ln(D)	-0.7714	0.9513	—	—	—
422 a+b·ln(D)	-5.613	2.5804	—	—	—
423 a+b·ln(D)	-7.47	1.6975	—	—	—
424 a·D <sup>b</sup> ·H <sup>c</sup>	0.1179	2.1052	-0.7931	—	—
425 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	37.78194	-16.7693	49	—	—
426 a·D <sup>2</sup>	37.78194	—	—	—	—
427 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	7.88144	19.3471	49	—	—
428 a·D <sup>2</sup>	7.88144	—	—	—	—
429 a+b·ln(D)	-5.478	2.494	—	—	—
430 a+b·ln(D)	-6.193	2.869	—	—	—
431 a+b·ln(D)	-6.621	3.155	—	—	—
432 a+b·H <sup>2</sup> +c·H <sup>3</sup>	0.4365	0.0033	0.0001	—	—
433 a+b·ln(D)	-5.777	2.481	—	—	—
434 a+b·[D/(D+7)]	-3.7983	7.7681	—	—	—
435 a+b·[D/(D+7)]+c·H+d·ln(H)	-3.4781	12.1095	0.0413	-1.5650	—
436 exp(a)·(D+1) <sup>b</sup> ·H <sup>c</sup> ·exp(d·D)·exp(e·H)	-4.634	4.496	-2.439	-0.0977	0.0984
437 a+b·ln(D)+c·ln(D <sup>2</sup> )+d·ln(H)	7.5174	1.29	0.4486	-1.1229	—
438 a+b·log(π·D)	-0.502	2.527	—	—	—
439 a·D <sup>b</sup>	0.00394	2.1534	—	—	—
440 a·D <sup>b</sup>	0.00083	2.4074	—	—	—
441 a·D <sup>b</sup>	0.33989	1.4728	—	—	—
442 a·D <sup>b</sup>	0.02157	2.221205	—	—	—
443 a+b·ln(D)	-3.636	1.977	—	—	—
444 a+b·ln(D)	-3.782	2.066	—	—	—
445 a+b·ln(D)	-4.079	2.252	—	—	—

## App. A

		Unit of Biom.	D	H	Range of D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	$r^2$
446 Poland	ln(RC)	kg	cm	—	—	—	62	8	15	0.909	
447 Sweden	ln(RC)	kg	cm	—	0–45	0—	50	8	26	305	0.901
448 Europe	RF	kg	cm	—	4.1–45	—	77	1	35	—	0.75
449 Sweden	ln(RS)	kg	cm	—	0–45	0—	50	8	25	286	0.889
450 —	log(RT)	kg	cm	—	—	—	68	8	—	—	0.965
451 —	log(RT)	kg	cm	—	—	—	68	8	17	—	—
452 —	log(RT)	kg	cm	m	—	—	68	8	6	0.966	
453 Europe	RT	kg	cm	—	4.1–45	—	77	1	35	—	0.964
454 Finland	log(RT)	kg	cm	—	7.0–21.6	7.9–14.8	56	8	5	—	0.935
455 Finland	log(RT)	kg	cm	—	4–24	—	21	8	20	0.99	
456 UK	RT	t	cm	—	—	—	9	6	—	—	—
457 Finland	log(SB)	kg	cm	m	5.6–23.6	8.0–18.2	56	8	—	0.968	
458 Finland	log(SB)	kg	cm	m	2.4–9.7	3.0–7.8	56	8	—	0.964	
459 Finland	log(SB)	kg	cm	m	7.0–21.6	7.9–14.8	56	8	—	0.961	
460 Finland	ln(SB)	kg	cm	m	5.1–45	3.5–27.5	40	8	—	0.981	
461 Finland	ln(SB)	kg	cm	—	—	—	55	8	18	0.98	
462 Finland	ln(SB)	kg	cm	—	—	—	55	8	30	0.95	
463 Norway	SB	gr	cm	—	7–15	—	6	7	16	0.999	
464 Norway	SB	gr	cm	—	2–7	—	6	7	16	—	
465 Norway	SB	gr	cm	—	7–20	—	6	7	30	16	0.993
466 Norway	SB	gr	cm	—	2–7	—	6	7	30	16	—
467 Sweden	ln(SB)	kg	cm	—	0–45	0—	50	8	461	0.929	
468 Sweden	ln(SB)	kg	cm	m	0–45	0—	50	8	461	0.935	
469 Sweden	SB	kg	cm	m	2–40	2—	1	8	73	0.984	
470 Sweden	ln(SB)	g	cm	dm	1.1–10	1.7–8.8	19	8	84	0.96	
471 UK	log(SB)	g	cm	—	—	—	48	8	10	0.98	
472 Sweden	SR	kg	cm	m	2–40	2—	1	8	73	0.943	
473 Sweden	SR	kg	cm	m	2–40	2—	1	8	73	0.962	
474 Sweden	ln(SR)	kg	cm	m	0–45	0—	50	8	296	0.958	
475 Belgium	ST	kg	cm	—	—	—	75	13	—	—	
476 Czech republic	ST	kg	cm	m	2–6	2–5	17	7	29	—	
477 Czech republic	ST	kg	cm	m	2–16	4–11	17	7	50	—	
478 Czech republic	ST	kg	cm	—	2–6	2–5	17	8	29	0.95	
479 Czech republic	ST	kg	cm	—	2–6	2–5	17	8	29	0.92	
480 Czech republic	ST	kg	cm	—	2–6	2–5	17	8	29	0.95	
481 Czech republic	ST	kg	cm	m	2–6	2–5	17	8	29	0.77	
482 Czech republic	ST	kg	cm	m	2–6	2–5	17	8	29	0.61	
483 Czech republic	ST	kg	cm	m	2–6	2–5	17	8	29	0.61	
484 Czech republic	ST	kg	cm	—	2–16	4–11	17	8	50	0.93	
485 Czech republic	ST	kg	cm	—	2–16	4–11	17	8	50	0.84	
486 Czech republic	ST	kg	cm	—	2–16	4–11	17	8	50	0.98	
487 Czech republic	ST	kg	cm	m	2–16	4–11	17	8	50	0.58	
488 Czech republic	ST	kg	cm	m	2–16	4–11	17	8	50	0.86	
489 Czech republic	ST	kg	cm	m	2–16	4–11	17	8	50	0.85	
490 Sweden	ln(ST)	kg	cm	—	0–45	0—	50	8	488	0.978	
491 Sweden	ln(ST)	kg	cm	m	0–45	0—	50	8	488	0.99	
492 Sweden	ST	kg	cm	m	2–40	2—	1	8	73	0.996	
493 UK	log(ST)	g	cm	—	—	—	48	8	10	0.995	
494 UK	log(ST)	g	cm	cm	—	—	63	8	—	—	
495 UK	log(ST)	g	cm	—	—	—	63	8	—	—	
496 UK	log(ST)	g	cm	cm	—	—	63	8	—	—	
497 UK	log(ST)	g	cm	—	—	—	63	8	—	—	
498 Finland	log(SU)	kg	cm	—	7–21.6	7.9–14.8	56	8	—	0.85	
499 Sweden	ln(SU)	kg	cm	—	0–45	0—	50	8	306	0.945	
500 Finland	log(SW)	kg	cm	m	5.6–23.6	8–18.2	56	8	—	0.994	
501 Finland	log(SW)	kg	cm	m	2.4–9.7	3–7.8	56	8	—	0.992	
502 Finland	log(SW)	kg	cm	m	7–21.6	7.9–14.8	56	8	—	0.991	
503 Finland	ln(SW)	kg	cm	m	121—	—	40	8	—	0.992	
504 Finland	SW	kg	cm	—	—	—	55	8	28	18	—
505 Finland	SW	kg	cm	—	—	—	55	8	29	30	—

Equation	a	b	Parameters		
			c	d	e
446 a+b·ln(D)	-3.514	1.909	-	-	-
447 a+b·[D/(D+10)]	-3.8375	8.8795	-	-	-
448 a·D <sup>b</sup>	0.038386	1.240689	-	-	-
449 a+b·[D/(D+9)]	-6.3413	13.2902	-	-	-
450 a·log(D)+b	2.2419	-1.3705	-	-	-
451 a·log(D)+b	2.6	-1.61	-	-	-
452 a·log(H·D <sup>2</sup> )+b	0.7665	-1.3736	-	-	-
453 a·D <sup>b</sup>	0.038872	2.066783	-	-	-
454 a+b·log(D)	-1.967	2.458	-	-	-
455 a+b·log(D)	-1.89	2.74	-	-	-
456 a·D <sup>b</sup>	5.595·10 <sup>-5</sup>	2.10019503	-	-	-
457 a+b·log(D <sup>2</sup> ·H)	-2.024	0.7603	-	-	-
458 a+b·log(D <sup>2</sup> ·H)	-1.827	0.7183	-	-	-
459 a+b·log(D <sup>2</sup> ·H)	-1.771	0.7106	-	-	-
460 a+b·ln(D <sup>2</sup> )+c·ln(H)	-4.344	0.885	0.435	-	-
461 a+b·ln(D)	-4.6637	2.4282	-	-	-
462 a+b·ln(D)	-3.5682	1.9976	-	-	-
463 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	22.63177	-6.7506	49	-	-
464 a·D <sup>2</sup>	22.63177	-	-	-	-
465 a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	23.31516	-6.2686	49	-	-
466 a·D <sup>2</sup>	23.31516	-	-	-	-
467 a+b·[D/(D+16)]	-2.9748	8.8489	-	-	-
468 a+b·[D/(D+16)]+c·ln(H)	-3.2765	7.2482	0.4487	-	-
469 exp(a)·(D+1) <sup>b</sup> ·exp(c·D)·exp(d·H)	-4.494	2.026	-0.0187	0.0494	-
470 a+b·ln(D)+c·ln(D <sup>2</sup> )+d·H	3.6931	0.9434	0.1778	0.0123	-
471 a+b·log(π·D)	0.007	2.146	-	-	-
472 exp(a)·D <sup>b</sup>	-6.158	3.071	-	-	-
473 exp(a)·D <sup>b</sup> ·H <sup>c</sup>	-6.383	2.613	0.569	-	-
474 a+b·[D/(D+12)]	-3.3913	11.1106	-	-	-
475 a·D <sup>b</sup>	0.12269	2.3272	-	-	-
476 a·(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.0038	4.4553	-1.5591	0.5661	-
477 a·(D+1) <sup>[b+c·log(D)]</sup> ·H <sup>d</sup>	0.006	3.1093	-0.5812	0.897	-
478 a+b·D	-1.666	0.928	-	-	-
479 a-exp(D·b)	0.1259	0.6526	-	-	-
480 a·D <sup>b</sup>	0.0867	2.2099	-	-	-
481 a+b·H	-1.572	0.8417	-	-	-
482 a-exp(H·b)	0.1348	0.5912	-	-	-
483 a·H <sup>b</sup>	0.0778	2.1498	-	-	-
484 a+b·D	-13.3349	3.223	-	-	-
485 a-exp(D·b)	1.0594	0.274	-	-	-
486 a·D <sup>b</sup>	0.1163	2.1826	-	-	-
487 a+b·H	-31.7127	5.3844	-	-	-
488 a-exp(H·b)	0.0814	0.584	-	-	-
489 a·H <sup>b</sup>	0.0025	3.9472	-	-	-
490 a+b·[D/(D+13)]	-2.3388	11.3264	-	-	-
491 a+b·[D/(D+13)]+c·H+d·ln(H)	-2.6768	7.5939	0.0151	0.8799	-
492 exp(a)·(D+1) <sup>b</sup> ·H <sup>c</sup> ·exp(d·H)	-3.760	1.882	0.758	0.0355	-
493 a+b·log(π·D)	0.981	2.194	-	-	-
494 a+b·log(H)	-1.34	1.89	-	-	-
495 a+b·log(D)	1.38	2.41	-	-	-
496 a+b·log(H)	-1.23	1.92	-	-	-
497 a+b·log(D)	1.4	2.64	-	-	-
498 a+b·log(D)	-1.740	0.9247	-	-	-
499 a+b·[D/(D+15)]	-3.9657	11.0481	-	-	-
500 a+b·log(D <sup>2</sup> ·H)	-1.730	0.9626	-	-	-
501 a+b·log(D <sup>2</sup> ·H)	-1.411	0.8316	-	-	-
502 a+b·log(D <sup>2</sup> ·H)	-1.663	0.9447	-	-	-
503 a+b·ln(D <sup>2</sup> )+c·ln(H)	-4.182	0.879	1.215	-	-
504 exp[a+b·ln(D)]+exp[c+d·ln(D)]	-3.9212	2.668	-13.575	5.2043	-
505 exp[a+b·ln(D)]+exp[c+d·ln(D)]	-2.5809	2.1595	-5.6832	2.407	-

## App. A

		Unit of Biom.	D	H	Range of D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	$r^2$
506 Norway	SW	gr	cm	—	7–15	—	6	7	16	0.992	
507 Norway	SW	gr	cm	—	2–7	—	6	7	16	—	
508 Norway	SW	gr	cm	—	7–20	—	6	7	30	16	0.991
509 Norway	SW	gr	cm	—	2–7	—	6	7	30	16	—
510 Sweden	ln(SW)	kg	cm	—	0–45	0—	50	8	461	0.966	
511 Sweden	ln(SW)	kg	cm	m	0–45	0—	50	8	461	0.986	
512 Sweden	ln(SW)	g	cm	dm	1.1–10	1.7–8.8	19	8	84	0.98	
<i>Pinus taeda</i>											
513 —	log(RT)	kg	cm	—	—	—	68	8	7	0.863	
<i>Populus tremula</i> (European aspen, Asp)											
514 Germany	AB	kg	cm	—	13.2–33	15.9–24.7	66	11	16	0.941	
515 Sweden	AB	kg	mm	—	1.9–9.2	3.6–15.8	36	4	—	0.958	
516 Sweden	BR	kg	mm	—	1.9–9.2	3.6–15.8	36	4	—	0.959	
517 Germany	CR	kg	cm	—	13.2–33	15.9–24.7	66	11	16	0.925	
518 Sweden	FL	kg	mm	—	1.9–9.2	3.6–15.8	36	4	—	0.944	
519 Germany	ST	kg	cm	—	13.2–33	15.9–24.7	66	11	16	0.909	
520 Sweden	ST	kg	mm	—	1.9–9.2	3.6–15.8	36	4	—	0.95	
<i>Populus trichocarpa</i> (Black cottonwood)											
521 Iceland	AB	kg	cm	m	4.6–34	4.6–20.7	71	8	22	0.98	
522 Iceland	CR	kg	cm	m	4.6–34	4.6–20.7	71	8	22	0.884	
523 Iceland	ST	kg	cm	m	4.6–34	4.6–20.7	71	8	22	0.98	
<i>Pseudotsuga menziesii</i> (Douglas-fir)											
524 Italy	ln(AB)	kg	cm	—	8.7–27	—	52	14	36	69	0.94
525 Netherlands	AB	kg	cm	—	3–38	6.7–25.9	4	14	23	0.995	
526 Netherlands	AB	kg	cm	—	5—	—	30	9	—	—	
527 Netherlands	ln(ABW)	kg	cm	—	5—	—	30	9	—	—	
528 Netherlands	BR	kg	cm	m	3–38	6.7–25.9	4	14	23	0.944	
529 Netherlands	CR	kg	cm	m	3–38	6.7–25.9	4	14	23	0.945	
530 UK	CR	kg	cm	m	—	—	9	6	2	—	—
531 Netherlands	FL	kg	cm	m	3–38	6.7–25.9	4	14	23	0.941	
532 —	log(RT)	kg	cm	—	—	—	68	8	18	0.908	
533 —	log(RT)	kg	cm	—	—	—	68	8	33	0.902	
534 —	log(RT)	kg	cm	—	—	—	68	8	14	0.907	
535 —	log(RT)	kg	cm	—	—	—	68	8	3	0.966	
536 —	log(RT)	kg	cm	m	—	—	68	8	3	0.947	
537 UK	RT	t	cm	—	—	—	9	6	—	—	
538 Netherlands	ST	kg	cm	m	3–38	6.7–25.9	4	14	23	0.998	
<i>Pseudotsuga</i> spp.											
539 Netherlands	log(RT)	kg	cm	—	5–27	—	21	8	21	0.96	
<i>Quercus conferta</i> (Hungarian oak, Platifolos dris)											
540 Greece	ln(ABW)	kg	cm	—	2–19	2.2–14.7	51	14	27	0.98	
541 Greece	ln(ABW)	kg	cm	cm	2–19	2.2–14.7	51	14	27	0.98	
542 Greece	ln(BR)	kg	cm	—	2–19	2.2–14.7	51	14	37	27	0.82
543 Greece	BR	kg	cm	—	2–19	2.2–14.7	51	14	38	27	0.65
544 Greece	ln(BR)	kg	cm	—	2–19	2.2–14.7	51	14	39	27	0.73
545 Greece	ln(BR)	kg	cm	—	2–19	2.2–14.7	51	14	40	27	0.89
546 Greece	ln(BR)	kg	cm	cm	2–19	2.2–14.7	51	14	37	27	0.79
547 Greece	BR	kg	cm	cm	2–19	2.2–14.7	51	14	38	27	0.5
548 Greece	ln(BR)	kg	cm	cm	2–19	2.2–14.7	51	14	39	27	0.68
549 Greece	ln(BR)	kg	cm	cm	2–19	2.2–14.7	51	14	40	27	0.86
550 Greece	ln(SW)	kg	cm	—	2–19	2.2–14.7	51	14	41	27	0.96
551 Greece	ln(SW)	kg	cm	—	2–19	2.2–14.7	51	14	42	27	0.89
552 Greece	ln(SW)	kg	cm	—	2–19	2.2–14.7	51	14	43	27	0.97
553 Greece	ln(SW)	kg	cm	cm	2–19	2.2–14.7	51	14	41	27	0.98
554 Greece	ln(SW)	kg	cm	cm	2–19	2.2–14.7	51	14	42	27	0.89
555 Greece	ln(SW)	kg	cm	cm	2–19	2.2–14.7	51	14	43	27	0.98
<i>Quercus ilex</i> (Holm oak)											
556 Italy	AB	kg	cm	—	20–90	—	73	14	—	—	
557 Italy	AB	kg	cm	—	5–20	—	47	14	12	—	
558 Italy	AB	kg	cm	m	4.5–26.1	6–16	7	8	94	0.952	

	Equation	a	b	Parameters		
				c	d	e
506	a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	89.24474	64.8925	49	—	—
507	a·D <sup>2</sup>	89.24474	—	—	—	—
508	a·D <sup>2</sup> +b·(D <sup>2</sup> -c)	175.40886	1.7509	49	—	—
509	a·D <sup>2</sup>	175.40886	—	—	—	—
510	a+b·[D/(D+14)]	-2.2184	11.4219	—	—	—
511	a+b·[D/(D+14)]+c·H+d·ln(H)	-2.6864	7.6066	0.0200	0.8658	—
512	a+b·ln(D)+c·ln(D <sup>2</sup> )+d·H	4.3823	1.19	0.1969	0.0149	—
513	a·log(D)+b	3.0742	-2.6683	—	—	—
514	a·D <sup>b</sup>	0.0519	2.545	—	—	—
515	a·D <sup>b</sup>	1.46·10 <sup>-4</sup>	2.6035333	—	—	—
516	a·D <sup>b</sup>	5.75·10 <sup>-4</sup>	1.873298	—	—	—
517	a·D <sup>b</sup>	0.0644	2.001	—	—	—
518	a·D <sup>b</sup>	8.47·10 <sup>-4</sup>	1.481578	—	—	—
519	a·D <sup>b</sup>	0.0197	2.764	—	—	—
520	a·D <sup>b</sup>	6.5·10 <sup>-5</sup>	2.739823	—	—	—
521	a·D <sup>b</sup> ·H <sup>c</sup>	0.0717	1.8322	0.6397	—	—
522	a·D <sup>b</sup> ·H <sup>c</sup>	0.0586	2.8285	-1.0282	—	—
523	a·D <sup>b</sup> ·H <sup>c</sup>	0.0379	1.581	1.0795	—	—
524	a+b·ln(D)	-1.957	2.2996	—	—	—
525	a+b·ln(D)	-1.62	2.410	—	—	—
526	a·D <sup>b</sup>	0.111	2.397	—	—	—
527	a·D <sup>b</sup>	0.111	2.397	—	—	—
528	a+b·ln(D)+c·ln(H)	-2.675	4.420	-2.784	—	—
529	a+b·ln(D)+c·ln(H)	-1.345	3.924	-2.514	—	—
530	a·D <sup>b</sup> ·H <sup>c</sup>	3.461·10 <sup>-4</sup>	2.7169	-1.26060	—	—
531	a+b·ln(D)+c·ln(H)	-1.346	3.351	-2.201	—	—
532	a·log(D)+b	2.1641	-1.4467	—	—	—
533	a·log(D)+b	2.5786	-1.8899	—	—	—
534	a·log(D)+b	2.9108	-2.3807	—	—	—
535	a·log(D)+b	2.5309	-1.6393	—	—	—
536	a·log(H·D <sup>2</sup> )+b	1.0472	-2.6287	—	—	—
537	a·D <sup>b</sup>	2.179·10 <sup>-5</sup>	2.4209	—	—	—
538	a+b·ln(D)+c·ln(H)	-2.535	2.009	0.709	—	—
539	a+b·log(D)	-2	2.63	—	—	—
540	a+b·ln(D)	-2.1686	2.4407	—	—	—
541	a+b·ln(D <sup>2</sup> ·H)	-2.5259	0.8605	—	—	—
542	a+b·ln(D)	-3.3508	1.7235	—	—	—
543	[a+b(1/D)+c(1/D <sup>2</sup> )]·D <sup>2</sup>	0.0536	-0.3269	—	—	—
544	a+b·ln(D)	-11.433	4.9391	—	—	—
545	a+b·ln(D)	-4.1909	2.5403	—	—	—
546	a+b·ln(D <sup>2</sup> ·H)	-3.5363	0.5957	—	—	—
547	[a+b(1/D <sup>2</sup> ·H)]·D <sup>2</sup> ·H	0.0015	0.0402	—	—	—
548	a+b·ln(D <sup>2</sup> ·H)	-12.7333	1.8202	—	—	—
549	a+b·ln(D <sup>2</sup> ·H)	-4.4702	0.8791	—	—	—
550	a+b·ln(D)	-2.5518	2.3887	—	—	—
551	a+b·ln(D)	-3.8649	2.4261	—	—	—
552	a+b·ln(D)	-2.32	2.4147	—	—	—
553	a+b·ln(D <sup>2</sup> ·H)	-2.9275	0.8468	—	—	—
554	a+b·ln(D <sup>2</sup> ·H)	-4.2122	0.854	—	—	—
555	a+b·ln(D <sup>2</sup> ·H)	-2.6916	0.8546	—	—	—
556	a·D <sup>b</sup>	0.2306	2.2791	—	—	—
557	a·D <sup>b</sup>	0.2179	2.0513	—	—	—
558	a+b·D <sup>2</sup> ·H	-0.6165	0.03582	—	—	—

## App. A

		Biom.	Unit of D	H	Range of D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	$r^2$	
559	Spain	log(AB)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	69	0.908	
560	Spain	log(AB)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	33	0.91	
561	Spain	log(AB)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	30	0.927	
562	Spain	log(AB)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	41	0.939	
563	Spain	log(AB)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	28	0.867	
564	Spain	log(AB)	kg	cm	m	5.3–24.4	4.4–12.8	12	8	63	0.942	
565	Spain	AB	kg	cm	—	—	—	25	14	15	—	
566	Italy	ABW	kg	cm	m	4.5–26.1	6–16	7	8	3	0.952	
567	Spain	log(BR)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	44	0.803	
568	Spain	log(BR)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	44	0.784	
569	Spain	log(BR)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	44	0.697	
570	Spain	log(BR)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	44	0.84	
571	Spain	log(BR)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	44	0.775	
572	Spain	log(BR)	kg	cm	m	5.3–24.4	4.4–12.8	12	8	44	0.807	
573	Spain	log(BR)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	45	0.781	
574	Spain	log(BR)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	46	0.443	
575	Italy	CR	kg	cm	m	4.5–26.1	6–16	7	8	94	0.952	
576	Spain	log(FL)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	69	0.615	
577	Spain	log(FL)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	33	0.467	
578	Spain	log(FL)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	30	0.65	
579	Spain	log(FL)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	41	0.735	
580	Spain	log(FL)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	28	0.74	
581	Spain	log(FL)	kg	cm	m	5.3–24.4	4.4–12.8	12	8	63	0.593	
582	Spain	log(RC)	kg	cm	—	7.2–23.1	4.4–12.8	13	8	25	0.64	
583	Spain	log(RC)	kg	cm	—	7.2–23.1	4.4–12.8	13	8	25	0.7	
584	Spain	log(RC)	kg	cm	—	8.8–19.8	5.5–9.0	13	8	25	0.7	
585	Spain	log(RS)	kg	cm	—	7.2–23.1	4.4–12.8	13	8	47	0.67	
586	Spain	log(RS)	kg	cm	—	7.2–23.1	4.4–12.8	13	8	47	0.75	
587	Spain	log(RS)	kg	cm	—	8.8–19.8	5.5–9.0	13	8	47	0.63	
588	Spain	log(RT)	kg	cm	—	7–23	—	21	8	32	0.73	
589	Spain	log(RT)	kg	cm	—	7.2–23.1	4.4–12.8	13	8	32	0.73	
590	Spain	log(RT)	kg	cm	—	7.2–23.1	4.4–12.8	13	8	20	0.81	
591	Spain	log(RT)	kg	cm	—	8.8–19.8	5.5–9.0	13	8	12	0.71	
592	Spain	log(ST)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	2	71	0.918
593	Spain	log(ST)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	2	33	0.929
594	Spain	log(ST)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	2	32	0.958
595	Spain	log(ST)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	2	43	0.941
596	Spain	log(ST)	kg	cm	—	5.3–24.4	4.4–12.8	12	8	2	28	0.882
597	Spain	log(ST)	kg	cm	m	5.3–24.4	4.4–12.8	12	8	2	65	0.963
<i>Quercus petraea</i>												
598	France	log(RT)	kg	cm	—	7–17	—	21	8	71	0.94	
<i>Quercus pyrenaica</i> (Pyrenean oak)												
599	Portugal	ln(SW)	kg	cm	m	2.5–46	3.3–27	14	8	166	0.99	
<i>Quercus</i> spp. (Oak, Eiche)												
600	Austria	ln(AB)	kg	cm	—	—	—	31	3	33	—	
601	UK	ln(ABW)	kg	cm	—	4.5–52	—	10	14	11	20	
602	UK	ln(ABW)	kg	cm	—	4.3–35	—	10	14	11	16	
603	UK	ln(ABW)	kg	cm	—	3.8–11	—	10	14	11	15	
604	UK	ln(ABW)	kg	cm	—	5.7–33	—	10	14	11	18	
605	UK	CR	t	cm	—	—	—	9	6	1	—	
606	UK	CR	t	cm	m	—	—	9	6	2	—	
<i>Tilia cordata</i> (Lime)												
607	UK	ln(ABW)	kg	cm	—	3.2–15	—	10	14	11	10	
											0.984	

Equation	a	b	Parameters		
			c	d	e
559 a+b·log(D)	-0.656	2.217	—	—	—
560 a+b·log(D)	-0.275	1.831	—	—	—
561 a+b·log(D)	-0.854	2.413	—	—	—
562 a+b·log(D)	-0.902	2.433	—	—	—
563 a+b·log(D)	-0.313	1.900	—	—	—
564 a+b·log(D)+c·H	-0.568	1.953	0.029	—	—
565 a·D <sup>b</sup>	0.2313	2.2662	—	—	—
566 a+b·D <sup>2</sup> ·H	-1.0906	0.031073	—	—	—
567 a+b·log(D)	-0.704	1.833	—	—	—
568 a+b·log(D)	-0.411	1.546	—	—	—
569 a+b·log(D)	-0.825	1.953	—	—	—
570 a+b·log(D)	-0.996	2.077	—	—	—
571 a+b·log(D)	-0.370	1.544	—	—	—
572 a+b·log(D)+c·H	-0.617	1.672	0.014	—	—
573 a+b·log(D)	-0.825	1.789	—	—	—
574 a+b·log(D)	-1.429	2.089	—	—	—
575 a+b·D <sup>2</sup> ·H	0.4741	0.0047473	—	—	—
576 a+b·log(D)	-1.624	1.891	—	—	—
577 a+b·log(D)	-1.347	1.654	—	—	—
578 a+b·log(D)	-2.128	2.309	—	—	—
579 a+b·log(D)	-2.142	2.269	—	—	—
580 a+b·log(D)	-1.366	1.774	—	—	—
581 a+b·log(D)+c·H	-1.533	1.808	0.002	—	—
582 a+b·log(D)	-1.188	2.139	—	—	—
583 a+b·log(D)	-1.563	2.422	—	—	—
584 a+b·log(D)	-0.714	1.79	—	—	—
585 a+b·log(D)	-1.145	1.897	—	—	—
586 a+b·log(D)	-1.417	2.093	—	—	—
587 a+b·log(D)	-0.835	1.688	—	—	—
588 a+b·log(D)	-1.05	2.19	—	—	—
589 a+b·log(D)	-1.047	2.191	—	—	—
590 a+b·log(D)	-1.393	2.451	—	—	—
591 a+b·log(D)	-0.448	1.734	—	—	—
592 a+b·log(D)	-1.166	2.478	—	—	—
593 a+b·log(D)	-0.747	2.044	—	—	—
594 a+b·log(D)	-1.355	2.674	—	—	—
595 a+b·log(D)	-1.336	2.640	—	—	—
596 a+b·log(D)	-0.839	2.156	—	—	—
597 a+b·log(D)+c·H	-1.088	2.157	0.039	—	—
598 a+b·log(D)	-1.56	2.44	—	—	—
599 a+b·ln(H·D <sup>2</sup> )	-3.323	0.95	—	—	—
600 a+b·ln(D)	-0.883	2.140	—	—	—
601 a+b·ln(D)	-2.4232	2.4682	—	—	—
602 a+b·ln(D)	-2.3223	2.4029	—	—	—
603 a+b·ln(D)	-3.1404	2.8113	—	—	—
604 a+b·ln(D)	-3.1009	2.6996	—	—	—
605 a·D <sup>2</sup>	$2.1612 \cdot 10^{-4}$	—	—	—	—
606 a·D <sup>b</sup> ·H <sup>c</sup>	$5.4224 \cdot 10^{-4}$	2.35	-1.022	—	—
607 a+b·ln(D)	-2.6788	2.4542	—	—	—

## Appendix A – References

- 1 Albrektson, A., Valinger, E. & Jonson, C. 1984. Några funktioner för bestämning av tallars biomassa i södra Norrland. Sveriges Skogsvårdsförbunds Tidskrift 82(6): 5–12.
- 2 Baldini, S., Berti, S., Cutini, A., Mannucci, M., Mercurio, R. & Spinelli, R. 1989. Prove sperimentali di primo diradamento in un soprassuolo di pino marittimo (*Pinus pinaster* Ait.) originato da incendio: aspetti silvicolturali, di utilizzazione e caratteristiche della biomassa. Ann. Ist. Sper. Selvicolatura 20: 385–436.
- 3 Barcikowski, A. & Loro, P.M. 1995. Needle biomass and dendrometrical features of Scots pine (*Pinus sylvestris* L.) natural regeneration seedlings of younger age classes, growing on a fresh coniferous forest site. Sylwan 139(2): 53–62.
- 4 Bartelink, H.H. 1996. Allometric relationships on biomass and needle area of Douglas-fir. Forest Ecology and Management 86: 193–203.
- 5 Bartelink, H.H. 1997. Allometric relationship for biomass and leaf area of beech (*Fagus sylvatica* L.). Annals of Forest Science 54: 39–50.
- 6 Brække, F.H. 1986. Distribution and yield of biomass from young *Pinus sylvestris* and *Picea abies* stands on drainages and fertilized peatland. Scandinavian Journal of Forest Research 1: 49–66.
- 7 Brandini, P. & Tabacchi, G. 1996. Biomass and volume equations for holm oak and strawberry-tree in coppice stands of Southern Sardinia. ISAFA Communicazioni di Ricerca 96(1): 59–69.
- 8 Briggs, E.F. & Cunia, T. 1982. Effect of cluster sampling in biomass tables construction: linear regression models. Canadian Journal of Forest Research 12(2): 255–263.
- 9 Broadmeadow, M. & Matthews, R. 2004. Survey methods for Kyoto Protocol monitoring and verification of UK forest carbon stocks. UK Emissions by Sources and Removals by Sinks due to Land Use, Land Use Change and Forestry Activities, Report. CEH, Edinburgh (June 2004).
- 10 Bunce, R.G.H. 1968. Biomass and production of trees in a mixed deciduous woodland. I. Girth and height as parameters for the estimation of tree dry weight. Journal of Ecology 56: 759–775.
- 11 Calamini, G. & Gregori, E. 2001. Study in a beech stand of Central Italy: allometric relations for the above ground biomass estimation. L'Italia Forestale e Montana 56(1): 1–23.
- 12 Canadell, J., Riba, M. & Andrés, P. 1988. Biomass equations for *Quercus ilex* L. in the Montseny Massif, Northeastern Spain. Forestry 61(2): 137–147.
- 13 Canadell, J. & Rodà, F. 1991. Root biomass of *Quercus ilex* in a montane mediterranean forest. Canadian Journal of Forest Research 21: 1771–1778.
- 14 Carvalho, J.P. & Parresol, B.R. 2003. Additivity in tree biomass components of Pyrenean oak (*Quercus pyrenaica* Willd.). Forest Ecology and Management 179: 269–276.
- 15 Cienciala, E., Cern\_, M., Alptauer, J. & Exnerová, Z. 2005. Biomass functions applicable to European beech. Journal of Forest Science 51(4): 147–154.
- 16 Cern\_, M. 1990. Biomass of *Picea abies* (L.) Karst. in midwestern Bohemia. Scandinavian Journal of Forest Research 5: 83–95.
- 17 Chroust, L. 1985. Above-ground biomass of young pine forests (*Pinus sylvestris*) and its determination. Communicationes Instituti Forestalis Cechosloveniae 14: 127–145.
- 18 Chroust, L. & Tesarova, J. 1985. Quantification of above-ground components of 20 years old Norway spruce (*Picea abies* (L.) Karsten). Communicationes Instituti Forestalis Cechosloveniae 14: 111–126.
- 19 Claesson, S., Sahlén, K. & Lundmark, T. 2001. Functions for biomass estimation of young *Pinus sylvestris*, *Picea abies* and *Betula* spp. from stands in northern Sweden with high stand densities. Scandinavian Journal of Forest Research 16: 138–146.
- 20 Dietrich, H.-P., Raspe, S., Schwarzmeier, M. & Ilg, S. 2002. Biomasse- und Nährrelementinventuren zur Ermittlung von Ernteeinzügen an drei bayerischen Fichtenstandorten. In: Dietrich, H.-P., Raspe, S. & Preuhlsler, T. (eds.): Inventur von Biomasse- und Nährstoffvorräten in Waldbeständen. Forstliche Forschungsberichte, München, 186: 59–72.
- 21 Drexhage, M. & Colin, F. 2001. Estimating root system biomass from breast-height diameters. Forestry 74(5): 491–497.
- 22 Drexhage, M. & Gruber, F. 1999. Above- and below-stump relationships for *Picea abies*: Estimating root system biomass from breast-height diameters. Scandinavian Journal of Forest Research 14: 328–333.
- 23 Duvigneaud, P., Kestemont, P., Timperman, J. & Moniquet, J.-C. 1977. La hêtre ardenne à *Festuca Altissima* a mirwart biomasse et productivité primaire. In: Duvigneaud, P. & Kestemont, P. (eds.). Productivité biologique en Belgique. Editions Duculot, Paris-Gembloux, p. 107–154.
- 24 Duvigneaud, P. & Timperman, P. 1977. Biomasse des épiphytes cryptogamiques dans une hêtre ardenne. In: Duvigneaud, P. & Kestemont, P. (eds.). Productivité biologique en Belgique. Editions Duculot, Paris-Gembloux.
- 25 Ferres, L., Roda, F., Verdu, A.M.C. & Terradas, J. 1980. Estructura y funcionalismo de un encinar montano en el Montseny. II. Biomasa aera. Mediterranea 4: 23–36.
- 26 Fiedler, F. 1986. Die Dendromasse eines hiefsreifen Fichtenbestandes. Beiträge für die Forstwirtschaft 20(4): 171–180.
- 27 Fiedler, F. 1987. Das Ökologische Meßfeld der Sektion Forstwirtschaft der TU Dresden. V. Die Verteilung der Dendromassekomponenten. Wissenschaftliche Zeitschrift der Technischen Universität Dresden 36(6): 229–234.

- 28 Hakkila, P. 1971. Coniferous branches as a raw material source. *Communicationes Instituti Forestalis Fenniae* 75(1): 1–60.
- 29 Hakkila, P. 1991. Hakkuupoistuman latvusmassa. *Folia Forestalia* 773: 1–24.
- 30 Hees, A.F.M.v. 2001. Biomass development in unmanaged forests. *Nederlands Bosbouwtijdschrift* 73(5): 2–5.
- 31 Hochbichler, E. 2002. Vorläufige Ergebnisse von Biomasseninventuren in Buchen- und Mittelwaldbeständen. In: Dietrich, H.-P., Raspe, S. & Preuhlsler, T. (eds.): Inventur von Biomasse- und Nährstoffvorräten in Waldbeständen. Forstliche Forschungsberichte, München, 186: 37–46.
- 32 Hughes, M.K. 1971. Tree biocontent, net production and litter fall in a deciduous woodland. *Oikos* 22: 62–73.
- 33 Ilg, S. 2002. Erhebung der oberirdischen Biomasse- und Elementvorräte eines Fichtenbestandes (*Picea abies* (L.) Karst.) im Bereich der Waldklimastation Zusmarshausen. Fachhochschule Weihenstephan. Series Erhebung der oberirdischen Biomasse- und Elementvorräte eines Fichtenbestandes (*Picea abies* (L.) Karst.) im Bereich der Waldklimastation Zusmarshausen. Fachbereich Forstwirtschaft. 98 p.
- 34 Ingerslev, M. & Hallbäcken, L. 1999. Above ground biomass and nutrient distribution in a limed and fertilized Norway spruce (*Picea abies*) plantation. Part II. Accumulation of biomass and nutrients. *Forest Ecology and Management* 119: 21–38.
- 35 Johansson, T. 1999a. Biomass equations for determining functions of pendula and pubescent birches growing on abandoned farmland and some practical implications. *Biomass and Bioenergy* 16: 223–238.
- 36 Johansson, T. 1999b. Biomass equations for determining functions of European aspen growing on abandoned farmland and some practical implications. *Biomass and Bioenergy* 17: 471–480.
- 37 Johansson, T. 1999c. Biomass production of Norway spruce (*Picea abies* (L.) Karst.) growing on abandoned farmland. *Silva Fennica* 33(4): 261–280.
- 38 Johansson, T. 1999d. Dry matter amounts and increment in 21- to 91-year-old common alder and grey alder and some practical implications. *Canadian Journal of Forest Research* 29: 1679–1690.
- 39 Johansson, T. 2000. Biomass equations for determining functions of common and grey alder growing on abandoned farmland and some practical implications. *Biomass and Bioenergy* 18: 147–159.
- 40 Korhonen, K.T. & Maltamo, M. 1990. Männyn maanpäällisten osien kuivamassat Etelä-Suomessa. *Metsätutkimustaitoksen tiedonantoja* 371: 1–29.
- 41 Laitat, E., Lebègue, C., Perrin, D. & Pissart, G. 2003. Séquestration du carbone par les forêts selon l'affectation des terres. Ministre de l'Agriculture et de la Ruralité, Ministère de la Région Wallonne, Division de la Nature et des Forêts, FuSAGxp.
- 42 Le Goff, N. & Ottorini, J.M. 2000. Biomass distributions at tree and stand levels in the beech experimental forest of Hesse (NE France). *Viterbo* (November): 9–11.
- 43 Le Goff, N. & Ottorini, J.M. 2001. Root biomass and biomass increment in a beech (*Fagus sylvatica* L.) stand in North-East France. *Annals of Forest Science* 58: 1–13.
- 44 Lee, D.-H. 1998. Architektur des Wurzelsystems von Fichten (*Picea abies* [L.] Karst.) auf unterschiedlich versauerter Standorten. Göttingen, Forschungszentrum Waldökosysteme der Universität Göttingen. Reihe A 153: 1–142.
- 45 Lee, D.-H. 2001. Relationship between above- and below-ground biomass for Norway spruce (*Picea abies*): estimating root system biomass from breast height diameter. *Jour. Korean For. Soc.* 90(3): 338–345.
- 46 Lehtonen, A. 2005. Estimating foliage biomass for Scots pine (*Pinus sylvestris* L.) and Norway spruce (*Picea abies* (L.) Karst.) plots. *Tree Physiology* 25: 803–811.
- 47 Leonardi, S. & Rapp, M. 1982. Phytomasse et mineralomase d'un taillis de Chene Vert du Massif de l'Etna. *Ecologia Mediterranea* 8: 125–138.
- 48 Lim, M.T. & J.E., C. 1986. The internal transfer of nutrients in a Scots pine stand: biomass components, current growth and their nutrient content. *Forestry* 59(1): 1–27.
- 49 Marklund, L.G. 1987. Biomass functions for Norway spruce (*Picea abies* (L.) Karst.) in Sweden. Sveriges lantbruksuniversitet, Institutionen för skogstaxering, Rapport 43: 1–127.
- 50 Marklund, L.G. 1988. Biomassafunktioner för tall, gran och björk i Sverige. Sveriges lantbruksuniversitet, Institutionen för skogstaxering, Rapport 45: 1–73.
- 51 Matis, K. & Alifragis, D. 1983–1984. Aboveground biomass of oaks (*Quercus conferta* Kit.) in Taxiarhis Greece. *Scientific Annals of the Depart. of Forestry and Natural Envir. KA* 15.
- 52 Menguzzato, G. & Tabacchi, G. 1986. Prove di diradamento su *Pseudotsuga menziesii* in Calabria. Ambiente, tavole di cubatura e della biomassa epigea. *Ann. Ist. Sper. Selvicoltura* 17: 255–293.
- 53 Menguzzato, G. 1988. Modelli di previsione del peso fresco, della biomassa e del volume per pino insigne ed eucalipti nell'Azienda Massanova (Salerno). *Ann. Ist. Sper. Selvicoltura* 19: 323–354.
- 54 Mund, M., Kummetz, E., Hein, M., Bauer, G.A. & Schulze, E.-D. 2002. Growth and carbon stocks of a spruce forest chronosequence in central Europe. *Forest Ecology and Management* 171: 275–296.
- 55 Mäkelä, A. & Vanninen, P. 1998. Impacts of size and competition on tree form and distribution of above-ground biomass in Scots pine. *Canadian Journal of Forest Research* 28: 216–227.
- 56 Mälkönen, E. 1974. Annual primary production and

- nutrient cycle in some Scots pine stands. *Communicationes Instituti Forestalis Fenniae* 84(5): 1–87.
- 57 Mälkönen, E. 1977. Annual primary production and nutrient cycle in birch stand. *Communicationes Instituti Forestalis Fenniae* 91(5): 1–35.
- 58 Mälkönen, E. & Saarsalmi, A. 1982. Hieskoivikon biomassatuotots ja ravinteiden menetys kokopuun korjuussa. *Folia Forestalia* 534: 1–17.
- 59 Møller, I.S. 2000. Calculation of biomass and nutrient removal for different harvesting intensities. *New Zealand Journal of Forestry Science* 30(1): 29–45.
- 60 Neumann, M. & Jandl, R. 2002. Aboveground biomass of two young Norway spruce stands. Manuscript.
- 61 Nihlgård, B. 1972. Plant biomass, primary production and distribution of chemical elements in a beech and a planted spruce forest in South Sweden. *Oikos* 23: 69–81.
- 62 Oleksyn, J., Reich, P.B., Chalupka, W. & Tjoelker, M.G. 1999. Differential above- and below-ground biomass accumulation of European *Pinus sylvestris* populations in a 12-year-old provenance experiment. *Scandinavian Journal of Forest Research* 14: 7–17.
- 63 Ovington, J.D. 1957. Dry matter production by *Pinus sylvestris* L. *Annals of Botany* 21(82): 287–314.
- 64 Poeppel, B. 1989. Untersuchungen der Dendromasse in mittelalten Fichtenbeständen. *Forsteinrichtung und Forstliche Ertragskunde*, Technische Universität Dresden. 66 p.
- 65 Pretzsch, H. 2000. Die Regeln von Reineke, Yoda und das Gesetz der räumlichen Allometrie. *Allgemeine Forst- und Jagd-Zeitung* 171: 205–210.
- 66 Rock, J. 2005. Suitability of published equations for aspen in Central Europe – results from a case study. Submitted manuscript.
- 67 Santa Regina, I. & Tarazona, T. 2001. Organic matter and nitrogen dynamics in a mature forest of common beech in the Sierra de la Demanda, Spain. *Ann. Sci. For.* 58: 301–314.
- 68 Santantonio, D., Hermann, R.K. & Overton, W.S. 1977. Root biomass studies in forest ecosystems. *Pedobiologia* 17: 1–31.
- 69 Schwarzmeier, M. 2000. Erhebung der oberirdischen Biomassevorräte von Fichtenbeständen (*Pinus abies* (L.) Karst.) im Bereich der Waldklimastationen Ebersberg und Flossenbürg. Diploma Thesis, Fachhochschule Weihenstephan, Fachbereich Forstwirtschaft, Germany 155 p.
- 70 Schöpfer, W. 1961. Beiträge zur Erfassung des Assimilationsapparates der Fichte. 127 p.
- 71 Snorrason, A. & Einarsson, S.F. 2004. Single-tree biomass and stem volume functions for eleven tree species used in Icelandic forestry. Submitted manuscript.
- 72 Starr, M., Hartman, M. & Kinnunen, T. 1998. Biomass functions for mountain birch in the Vuoskojärvi Integrated Monitoring area. *Boreal Environment Research* 3: 297–303.
- 73 Susmel, L., Viola, F. & Bassalo, G. 1976. Ecologie della Lecceta del Supramonte di Orgosolo. (Sardegna Centro-orientale). *Analisi de Centro di Economia Montana delle Venezie* 10: 1–216.
- 74 Wirth, C., Schumacher, J. & Schulze, E.-D. 2004. Generic biomass functions for Norway spruce in central Europe – a meta-analysis approach toward prediction and uncertainty estimation. *Tree Physiology* 24: 121–139.
- 75 Xiao, C.W., C.Y., J., Janssens, I.A., Roskams, P., Nachtergale, L., Carrara, A., Sanchez, B.Y. & Ceulemans, R. 2003. Above-and belowground biomass and net primary production in a 73-year-old Scots pine forest. Manuscript.
- 76 Zianis, D. & Mencuccini, M. 2003. Aboveground biomass relationship for beech (*Fagus moesiaca* Cz.) trees in Vermio Mountain, Northern Greece, and generalised equations for *Fagus* sp. *Annals of Forest Science* 60: 439–448.
- 77 The equation is derived from 3 different sources:
- Makkonen, K. & Helmisaari, H.-S. 1998. Seasonal and yearly variations of fine-root biomass and necromass in a Scots pine (*Pinus sylvestris* L.) stand. *Forest Ecology and Management* 102: 283–290.
- Oleksyn, J., Reich, P.B., Chalupka, W. & Tjoelker, M.G. 1999. Differential above- and below-ground biomass accumulation of European *Pinus sylvestris* populations in a 12-year-old provenance experiment. *Scandinavian Journal of Forest Research* 14: 7–17.
- Vanninen, P., Ylitalo, H., Sievänen, R. & Mäkelä, A. 1996. Effects of age and site quality on the distribution of biomass in Scots pine (*Pinus sylvestris* L.). *Trees* 10: 231–238.

## Appendix A – Contact persons

- Boosten M.**, Martijn.boosten@wur.nl, Wageningen University, the Netherlands
- Cienciala E.**, Emil.Cienciala@ifer.cz, Institute of Forest Ecosystem Research (IFER), Czech Republic
- Hochbichler E.**, ehochbic@edv1.boku.ac.at, Universität für Bodenkultur, Wien
- Johansson T.**, tord.johansson@bioenergi.slu.se, SLU, Sweden
- Laitat E.**, eric.laitat@cec.eu.int, Department of Plant Biology, Gembloux Agricultural university
- Matthews, R.**, robert.matthews@forestry.gsi.gov.uk, Forest Research, Alice Holt Research Station, UK
- Møller I.S.**, ism@fsl.dk, Danish centre for Forest, Landscape and Planning
- Muukkonen P.**, petteri.muukkonen@metla.fi, Finnish Forest Research Institute, Finland

- 9 **Nabuurs G.-J.**, G.J.Nabuurs@Alterra.wag-ur.nl, Alterra, Wageningen, the Netherlands
- 10 **Raspe S.**, ras@lwf.uni-muenchen.de, Bayerische Landesanstalt für Wald und Forstwirtschaft, Freising, Germany
- 11 **Rock, J.**, rock@pik-potsdam.de, Global Change and Natural Systems Department, Potsdam Institute for Climate Change Impact Research
- 12 **Wirth C.**, cwirth@bgc-jena.mpg.de, Max-Planck-Institute for Biogeochemistry, Jena, Germany
- 13 **Xiao C.W.**, Xiao@uia.ua.ac.be, University of Antwerpen, Belgium
- 14 **Zianis D.**, IERM, Edinburgh University, UK
- 25 Roots more than 5 cm in diameter
- 26 Roots less than 5 cm in diameter
- 28 Dominant trees over an age gradient
- 29 Trees of different sizes in one age group
- 30 Fertilised plot
- 31 Doubt about the parameters.
- 32 Doubt about the sign of the parameter c. The sign of the parameter c has been changed on this table.
- 33 One year old needles
- 34 Two years old needles
- 35 The equation is derived from 3 different sources:  
Makkonen, K. & Helmisaari, H.-S. 1998. Seasonal and yearly variations of fine-root biomass and necromass in a Scots pine (*Pinus sylvestris* L.) stand. *Forest Ecology and Management* 102: 283–290.  
Oleksyn, J., Reich, P.B., Chalupka, W. & Tjoelker, M.G. 1999. Differential above- and below-ground biomass accumulation of European *Pinus sylvestris* populations in a 12-year-old provenance experiment. *Scandinavian Journal of Forest Research* 14: 7–17.  
Vanninen, P., Ylitalo, H., Sievänen, R. & Mäkelä, A. 1996. Effects of age and site quality on the distribution of biomass in Scots pine (*Pinus sylvestris* L.). *Trees* 10: 231–238.
- The equation is based on 38 different values of total root biomass and diameter at breast height; some values were derived from individual trees; some values were averages from several trees.
- 36 Regressions developed by D Zianis. Original data from graphs
- 37 Branches less than or equal to 1 cm
- 38 Branches between 1 and 3 cm
- 39 Branches bigger than 3 cm
- 40 All branches
- 41 Stem wood beneath crown
- 42 Stem wood in crown
- 43 Total stem wood
- 44 Branches less than 5 cm in diameter
- 45 Branches 1 to 5 cm in diameter
- 46 Branches greater than 5 cm in diameter

## Appendix A – Comments

- 1 Trees 7 cm or less in diameter
- 2 Trees more than 7 cm in diameter
- 3 Merchantable parts, >3–4 cm
- 4 Living branches
- 5 Roots more than 1 cm in diameter
- 6 Regressions developed by D Zianis. Original data from tables
- 7 0.3 m from forest floor
- 8 Young trees
- 9 Middle age trees
- 10 Adult trees
- 11 Original regressions relate girth to biomass
- 12 D at tree base
- 13 Other types of functions with D and H and independent variables are given for the same material, includes determination coefficients
- 14 Newly cut trees
- 15 Cut trees dried in the stand during summer with a total loss of needles of about 75%
- 16 Includes fertilized plots
- 17 Wood of all living branches
- 18 Branch and top wood
- 19 Wood and bark of all living branches
- 20 Twigs
- 21 Living crown
- 22 Whole crown
- 23 Other than current year needles, includes fertilized plots
- 24 Includes needles, twigs, and branches less than 1 cm in diameter

**Appendix B.** General descriptions of volume equations. Both scientific and common names of the tree species are shown, since that helps in reading the original papers. Number of sampled trees ( $n$ ), coefficients of determination ( $r^2$ ), and range of diameter ( $D$ ) and height ( $H$ ) of sampled trees are reported when available in original article. References (Ref.) to the original papers according to author as well as the contact (Cont.) person who submitted the equation to this database are given at the end of the table. In the comments column (Comm.) occur some comments about the particular equation.

	Unit of Vol.	$D$	$H$	Range of $D$ (cm)	Range of $H$ (m)	Ref.	Cont.	Comm.	$n$	$r^2$
<i>Abies alba</i> (Silver fir)										
1 Norway	dm <sup>3</sup>	cm	m	5–	–	37	9	1	–	–
<i>Abies grandis</i> (Grand fir)										
2 Netherlands	dm <sup>3</sup>	cm	m	–	–	16	4		285	0.996
3 Norway	dm <sup>3</sup>	cm	m	5–	–	37	9	1	–	–
<i>Abies sibirica</i> (Fir)										
4 Germany	m <sup>3</sup>	cm	–	10–48	–	20	3		–	–
<i>Abies</i> spp. (Fir, Brad)										
5 Austria	dm <sup>3</sup>	dm	dm	1–	–	29	5		–	–
6 Austria	dm <sup>3</sup>	dm	dm	0.5–1.04	–	32	5		–	0.848
7 Romania	m <sup>3</sup>	cm	m	–	–	18	3		–	–
<i>Acacia</i> spp. (Salcim)										
8 Romania	m <sup>3</sup>	cm	m	–	–	18	3		–	–
<i>Acer pseudoplatanus</i> (Maple, Erable sycomore, Paltin, Sycamore)										
9 Belgium	m <sup>3</sup>	cm	m	–	–	14	6	2	419	–
10 Netherlands	dm <sup>3</sup>	cm	m	–	–	16	4		143	0.99
11 Romania	m <sup>3</sup>	cm	m	–	–	18	3		–	–
12 UK	m <sup>3</sup>	cm	m	–	–	8	2		–	–
<i>Alnus alba</i> (Anin alb)										
13 Romania	m <sup>3</sup>	cm	m	–	–	18	3		–	–
<i>Alnus glutinosa</i> (Black alder, Klibbal)										
14 Netherlands	dm <sup>3</sup>	cm	m	–	–	16	4		55	0.991
15 Norway	dm <sup>3</sup>	cm	m	10–	–	7	9		–	–
16 Norway	dm <sup>3</sup>	cm	m	0–12	–	7	9		–	–
17 Sweden	dm <sup>3</sup>	cm	m	–	–	17	3	3	1643	–
18 Sweden	dm <sup>3</sup>	cm	m	–	–	17	3	3	1643	–
19 Sweden	dm <sup>3</sup>	cm	m	–	–	17	3	4	1643	–
<i>Alnus incana</i> (Grey alder)										
20 Norway	dm <sup>3</sup>	cm	m	5–	–	3	9		–	–
<i>Alnus nigra</i> (Anin negru)										
21 Romania	m <sup>3</sup>	cm	m	–	–	18	3		–	–
<i>Alnus</i> spp. (Alder)										
22 Austria	dm <sup>3</sup>	dm	dm	0.5–1.04	–	32	5		–	0.583
<i>Arbutus unedo</i> (Strawberry-tree)										
23 Italy	dm <sup>3</sup>	cm	–	6–15	–	6	3	5	26	0.965
<i>Betula pendula</i> (Birch, Berk)										
24 Netherlands	dm <sup>3</sup>	cm	m	–	–	16	4		27	0.999
<i>Betula</i> spp. (Birch, Björk, Bjørk, Bouleaux, Mesteacan)										
25 Belgium	m <sup>3</sup>	cm	m	–	–	14	6	2	329	–
26 Finland	ln(dm <sup>3</sup> )cm	–	–	1.2–49.7	2.4–29.5	22	3		863	–
27 Finland	dm <sup>3</sup>	cm	–	1.2–49.7	2.4–29.5	22	3		863	–
28 Finland	dm <sup>3</sup>	cm	m	1.2–49.7	2.4–29.5	22	3		863	–
29 Finland	m <sup>3</sup>	cm	m	–	–	19	3		863	–
30 Norway	m <sup>3</sup>	cm	m	2–44	4–28	3	3		3312	0.984
31 Norway	m <sup>3</sup>	cm	m	2–44	4–28	3	3		722	0.979
32 Romania	m <sup>3</sup>	cm	m	–	–	18	3		–	–
33 Sweden	dm <sup>3</sup>	cm	m	5–34.9	5.0–26.9	27	3		1746	–
34 Sweden	dm <sup>3</sup>	cm	m	4.5–	6–	5	3	6	761	0.992
35 Sweden	dm <sup>3</sup>	cm	m	4.5–	6–	5	3	7	761	0.988
36 Sweden	dm <sup>3</sup>	cm	m	4.5–	6–	5	3	8	521	0.995
37 Sweden	dm <sup>3</sup>	cm	m	4.5–	6–	5	3	9	521	0.993

App. B	Unit of Vol.	D	H	Range of D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	$r^2$
38 Sweden	dm <sup>3</sup>	cm	m	5–34.9	5–26.9	27	3		1363	—
39 UK	m <sup>3</sup>	cm	m	—	—	8	2		—	—
<i>Carpinus</i> spp.										
40 Netherlands	dm <sup>3</sup>	mm	m	—	—	31	3	10	—	—
41 Netherlands	dm <sup>3</sup>	mm	m	—	—	31	3	11	—	—
42 Netherlands	dm <sup>3</sup>	mm	m	—	—	31	3	12	—	—
<i>Chamaecyparis lawsoniana</i> (Lawson cypress)										
43 Netherlands	dm <sup>3</sup>	cm	m	—	—	16	4		101	0.987
<i>Corylus avellana</i> (Hazel)										
44 Norway	dm <sup>3</sup>	cm	m	5–	—	3	9		—	—
<i>Fagus</i> spp. (Beech, Fag.)										
45 Austria	dm <sup>3</sup>	dm	dm	1–	—	29	5		—	—
46 Austria	dm <sup>3</sup>	dm	dm	0.5–1.04	—	32	5		—	0.748
47 Romania	m <sup>3</sup>	cm	m	—	—	18	3		—	—
48 UK	m <sup>3</sup>	cm	m	—	—	8	2		—	—
<i>Fagus sylvatica</i> (Beech, Rotbuche, Beuk)										
49 Belgium	m <sup>3</sup>	cm	m	—	—	14	6		—	—
50 Germany	dm <sup>3</sup>	cm	dm	10.7–61.8	10.2–34.6	28	1	13	20	0.994
51 Germany	dm <sup>3</sup>	cm	dm	10.7–61.8	10.2–34.6	28	1	14	20	0.973
52 Netherlands	dm <sup>3</sup>	cm	m	—	—	16	4		30	0.999
53 Netherlands	dm <sup>3</sup>	cm	m	—	—	15	3		—	—
<i>Fraxinus excelsior</i> (Ash, Frêne, Es)										
54 Belgium	m <sup>3</sup>	cm	m	—	—	14	6	2	534	—
55 Netherlands	dm <sup>3</sup>	cm	m	—	—	16	4		121	0.984
56 Sweden	dm <sup>3</sup>	cm	m	—	—	17	3	15	5294	—
57 Sweden	dm <sup>3</sup>	cm	m	—	—	17	3	16	5294	—
58 Sweden	dm <sup>3</sup>	cm	m	—	—	17	3	15	5294	—
59 Sweden	dm <sup>3</sup>	cm	m	—	—	17	3	17	5284	—
60 Sweden	dm <sup>3</sup>	cm	m	—	—	17	3	18	5284	—
<i>Fraxinus</i> spp. (Ash)										
61 Norway	dm <sup>3</sup>	cm	m	5–	—	3	9		—	—
62 Romania	m <sup>3</sup>	cm	m	—	—	18	3		—	—
63 UK	m <sup>3</sup>	cm	m	—	—	8	2		—	—
<i>Larix decidua</i> (Larch, Mélèzes)										
64 Austria	dm <sup>3</sup>	dm	dm	1–	—	29	5		—	—
65 Austria	dm <sup>3</sup>	dm	dm	0.5–1.04	—	32	5		—	0.775
66 Belgium	m <sup>3</sup>	cm	m	—	—	14	6	2	503	—
67 Netherlands	dm <sup>3</sup>	cm	m	—	—	16	4		123	0.996
68 Norway	dm <sup>3</sup>	cm	m	5–	—	37	9	1	—	—
<i>Larix hybrid</i> (Hybrid larch)										
69 Norway	dm <sup>3</sup>	cm	m	5–	—	37	9	1	—	—
<i>Larix kaempferi</i> (Japanese larch)										
70 Netherlands	dm <sup>3</sup>	cm	m	—	—	16	4		1023	0.996
71 Norway	dm <sup>3</sup>	cm	m	5–	—	37	9	1	—	—
<i>Larix sibirica</i> (Siberian larch)										
72 Iceland	m <sup>3</sup>	cm	m	4–34	4–16	26	7		100	0.993
73 Iceland	m <sup>3</sup>	cm	m	4–34	4–16	26	7	19	100	0.991
74 Iceland	m <sup>3</sup>	cm	m	3.3–31.6	3–20	33	3		44	0.995
75 Norway	dm <sup>3</sup>	cm	m	5–	—	37	9	1	—	—
<i>Larix</i> spp. (Lehtikuusi, Lork, Larice)										
76 Finland	dm <sup>3</sup>	cm	m	—	2–10	35	3		2813	—
77 Finland	dm <sup>3</sup>	cm	m	—	2–10	35	3		2813	—
78 Netherlands	dm <sup>3</sup>	mm	m	—	—	31	3	10	—	—
79 Netherlands	dm <sup>3</sup>	mm	m	—	—	31	3	11	—	—
80 Netherlands	dm <sup>3</sup>	mm	m	—	—	31	3	12	—	—
81 Romania	m <sup>3</sup>	cm	m	—	—	18	3		—	—
<i>Picea abies</i> (Norway spruce, Kuusi, Gran, Epicéa, Fijnspar)										
82 Austria	dm <sup>3</sup>	dm	dm	1–	—	29	5		—	—
83 Austria	dm <sup>3</sup>	dm	dm	0.5–1.04	—	32	5		—	0.812
84 Belgium	m <sup>3</sup>	cm	m	—	—	14	6	2	991	—

## App. B

	Unit of Vol.	D	H	Range of D (cm)	H (m)	Ref.	Cont.	Comm.	n	$r^2$
85 Czech Republic	$m^3$	cm	m	—	—	12	3	26	0.97	
86 Czech Republic	$m^3$	cm	m	—	—	12	3	26	0.98	
87 Finland	$\ln(dm^3)cm$	—	—	1.5–61.9	1.8–32.7	22	3	1864	—	
88 Finland	$\ln(dm^3)cm$	—	—	1.5–61.9	1.8–32.7	22	3	1864	—	
89 Finland	$dm^3$	cm	m	1.5–61.9	1.8–32.7	22	3	1864	—	
90 Finland	$\ln(dm^3)cm$	m	—	—	—	19	3	2864	—	
91 Finland	$dm^3$	cm	m	2–18	2–18	21	3	359	0.993	
92 Finland	$dm^3$	cm	m	2–18	2–18	21	3	180	0.995	
93 Finland	$\ln(dm^3)cm$	m	—	7–45	5–29	23	3	744	—	
94 Germany	$m^3$	m	m	4.9–10.3	5–8.8	25	3	5	0.976	
95 Germany	$m^3$	m	m	8.7–12.3	7.6–10.2	25	3	5	0.998	
96 Iceland	$m^3$	cm	m	2.7–27.9	2.7–12	33	3	16	0.994	
97 Netherlands	$dm^3$	mm	m	—	—	31	3	10	—	
98 Netherlands	$dm^3$	mm	m	—	—	31	3	11	—	
99 Netherlands	$dm^3$	mm	m	—	—	31	3	12	—	
100 Netherlands	$dm^3$	cm	m	—	—	16	4	745	0.995	
101 Netherlands	$dm^3$	cm	m	—	—	15	3	—	—	
102 Norway	$dm^3$	cm	m	—	—	1	3	20	2621	
103 Norway	$dm^3$	cm	m	—	—	1	3	20	1348	
104 Norway	$dm^3$	cm	m	—	—	1	3	21	1813	
105 Norway	$dm^3$	cm	m	—	—	34	3	3597	0.988	
106 Norway	$dm^3$	cm	m	—	—	34	3	—	—	
107 Norway	$dm^3$	cm	m	10.1–12.9	—	34	3	7446	0.988	
108 Norway	$dm^3$	cm	m	13–59.4	—	34	3	6096	0.988	
109 Norway	$dm^3$	cm	m	10–59.4	—	34	3	1350	0.993	
110 Norway	$dm^3$	cm	m	—	—	34	3	699	0.988	
111 Norway	$dm^3$	cm	m	10–59.4	—	34	3	2004	0.986	
112 Norway	$dm^3$	cm	m	—	—	34	3	2898	0.988	
113 Norway	$dm^3$	cm	m	10–59.4	—	34	3	5442	0.99	
114 Poland	$m^3$	cm	m	—	—	10	3	2000	—	
115 Poland	$m^3$	cm	m	—	—	9	3	22	2000	
116 Sweden	$dm^3$	cm	m	5–55.9	3–34.9	27	3	3925	—	
117 Sweden	$dm^3$	cm	m	—	—	4	3	2384	—	
118 Sweden	$dm^3$	cm	m	—	—	5	3	2609	0.998	
119 Sweden	$dm^3$	cm	m	—	—	5	3	6	2171	
120 Sweden	$dm^3$	cm	m	—	—	5	3	9	2609	
121 Sweden	$dm^3$	cm	m	—	—	5	3	7	2171	
122 Sweden	$dm^3$	cm	m	—	—	27	3	2424	—	
123 Sweden	$dm^3$	cm	m	—	—	4	3	2173	—	
<i>Picea engelmannii</i> (Engelmanni spruce)										
124 Iceland	$m^3$	cm	m	—	—	33	3	15	0.968	
<i>Picea sitchensis</i> (Sitka spruce)										
125 Netherlands	$dm^3$	cm	m	—	—	16	4	85	0.997	
126 Norway	$dm^3$	cm	m	—	—	1	3	20	2429	
127 Norway	$dm^3$	cm	m	—	—	1	3	20	1447	
128 Norway	$dm^3$	cm	m	—	—	1	3	21	1363	
<i>Picea</i> spp. (Molid)										
129 Romania	$m^3$	cm	m	—	—	18	3	—	—	
130 Iceland	$m^3$	cm	m	—	—	33	3	56	—	
<i>Pinus contorta</i> (Contorta tall)										
131 Iceland	$m^3$	cm	m	—	—	33	3	48	—	
132 Netherlands	$dm^3$	cm	m	—	—	16	4	127	0.994	
133 Sweden	$dm^3$	cm	m	—	—	17	3	3	1301	
134 Sweden	$dm^3$	cm	m	—	—	17	3	3	1301	
135 Sweden	$dm^3$	cm	m	—	—	17	3	22	1301	
<i>Pinus nigra</i> var <i>maritima</i> (Black pine)										
136 Netherlands	$dm^3$	cm	m	—	—	16	4	798	0.997	
<i>Pinus nigra</i> var <i>nigra</i> (Black pine, Pin negru)										
137 Netherlands	$dm^3$	cm	m	—	—	16	4	983	0.996	
138 Romania	$m^3$	cm	m	—	—	18	3	—	—	

App. B	Unit of Vol.	D	H	Range of D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	$r^2$
<b><i>Pinus</i> spp.</b>										
139 Germany	$m^3$	m	—	—	—	20	3	—	—	—
140 Germany	$m^3$	cm	—	5–75	—	20	3	—	—	—
141 Netherlands	$dm^3$	mm	m	—	—	31	3	10	—	—
142 Netherlands	$dm^3$	mm	m	—	—	31	3	11	—	—
143 Netherlands	$dm^3$	mm	m	—	—	31	3	12	—	—
<b><i>Pinus sylvestris</i> (Scots pine, Mänty, Tall, Furu, Grove den, Pin silvestri)</b>										
144 Austria	$dm^3$	dm	dm	0.5–	—	29	5	—	—	—
145 Belgium	$m^3$	cm	—	—	—	36	8	—	—	—
146 Belgium	$m^3$	cm	m	—	—	14	6	—	—	—
147 Finland	$ln(dm^3)$	cm	—	0.9–50.6	1.5–28.3	22	3	—	2050	—
148 Finland	$ln(dm^3)$	cm	—	0.9–50.6	1.5–28.3	22	3	—	2050	—
149 Finland	$dm^3$	cm	m	0.9–50.6	1.5–28.3	22	3	—	2050	—
150 Finland	$ln(dm^3)$	cm	m	—	—	19	3	—	2326	—
151 Finland	$dm^3$	cm	m	2–18	2–18	21	3	—	249	0.994
152 Finland	$dm^3$	cm	m	2–18	2–18	21	3	—	486	0.989
153 Finland	$dm^3$	cm	m	—	—	35	3	—	1493	—
154 Finland	$ln(dm^3)$	cm	m	7–50	5–28	23	3	—	1291	—
155 Germany	$m^3$	cm	m	3–14	5.8–10.7	24	3	—	—	—
156 Italy	$dm^3$	dm	dm	13–49	7–27.5	13	3	—	114	—
157 Netherlands	$dm^3$	mm	m	—	—	31	3	10	—	—
158 Netherlands	$dm^3$	mm	m	—	—	31	3	11	—	—
159 Netherlands	$dm^3$	mm	m	—	—	31	3	12	—	—
160 Netherlands	$dm^3$	cm	m	—	—	16	4	—	1207	0.994
161 Norway	$dm^3$	cm	m	—	—	1	3	20	4816	0.997
162 Norway	$dm^3$	cm	m	—	—	1	3	20	986	0.997
163 Norway	$dm^3$	cm	m	—	—	1	3	21	3010	0.994
164 Norway	$dm^3$	cm	m	10–	—	7	3	—	4356	0.996
165 Norway	$dm^3$	cm	m	—12	—	7	3	—	2622	0.999
166 Norway	$dm^3$	cm	m	—12	—	7	3	—	2622	0.999
167 Norway	$dm^3$	cm	m	—12	—	7	3	—	2622	0.999
168 Norway	$dm^3$	cm	m	—12	—	7	3	—	2622	0.999
169 Romania	$m^3$	cm	m	—	—	18	3	—	—	—
170 Sweden	$dm^3$	cm	m	5–49.9	3–32.9	27	3	—	4421	—
171 Sweden	$dm^3$	cm	m	2–	2–	4	3	—	3407	—
172 Sweden	$dm^3$	cm	m	4.5–	4–	5	3	8	3734	0.996
173 Sweden	$dm^3$	cm	m	4.5–	4–	5	3	6	2215	0.996
174 Sweden	$dm^3$	cm	m	4.5–	4–	5	3	9	3734	0.993
175 Sweden	$dm^3$	cm	m	4.5–	4–	5	3	7	2215	0.994
176 Sweden	$dm^3$	cm	m	5–49.9	3–32.9	27	3	—	2390	—
177 Sweden	$dm^3$	cm	m	2–	2–	4	3	—	2071	—
<b><i>Populus</i> spp. (Poplar, Plop)</b>										
178 Austria	$dm^3$	dm	dm	5–10.4	—	32	5	—	—	0.849
179 Netherlands	$dm^3$	mm	m	—	—	31	3	10	—	—
180 Netherlands	$dm^3$	mm	m	—	—	31	3	11	—	—
181 Netherlands	$dm^3$	mm	m	—	—	31	3	12	—	—
182 Romania	$m^3$	cm	m	—	—	18	3	—	—	—
183 Romania	$m^3$	cm	m	—	—	18	3	—	—	—
184 UK	$m^3$	cm	m	—	—	8	2	—	—	—
<b><i>Populus tremula</i> (Aspen, Plop tremulator)</b>										
185 Norway	$dm^3$	cm	m	13–	—	11	9	—	—	—
186 Norway	$dm^3$	cm	m	0–13	—	11	9	—	—	—
187 Romania	$m^3$	cm	m	—	—	18	3	—	—	—
188 Sweden	$dm^3$	cm	m	—	—	17	3	3	707	—
189 Sweden	$dm^3$	cm	m	—	—	17	3	3	707	—
190 Sweden	$dm^3$	cm	m	—	—	17	3	22	707	—
<b><i>Populus trichocarpa</i> (Black cottonwood)</b>										
191 Iceland	$m^3$	cm	m	4.6–34	4.6–20.7	33	3	—	25	0.989
<b><i>Prunus avium</i> (Wild cherry, Merisier, Zoete kers)</b>										
192 Belgium	$m^3$	cm	m	—	—	14	6	2	334	—

## App. B

	Unit of Vol.	D	H	Range of D (cm)	Range of H (m)	Ref.	Cont.	Comm.	n	$r^2$
<i>Pseudotsuga menziesii</i> (Douglas fir, Duglas)										
193 Belgium	$m^3$	cm	m	—	—	14	6	2	632	—
194 Netherlands	$dm^3$	cm	m	—	—	16	4		1136	0.993
195 Norway	$dm^3$	cm	m	5—	—	37	9	1	—	—
196 Romania	$m^3$	cm	m	—	—	18	3		—	—
<i>Pseudotsuga</i> spp.										
197 Netherlands	$dm^3$	mm	m	—	—	31	3	10	—	—
198 Netherlands	$dm^3$	mm	m	—	—	31	3	11	—	—
199 Netherlands	$dm^3$	mm	m	—	—	31	3	12	—	—
<i>Quercus grisea</i> (Gray oak, Stejar brumariu)										
200 Romania	$m^3$	cm	m	—	—	18	3		—	—
<i>Quercus ilex</i> (Holm oak)										
201 Italy	$dm^3$	cm	m	4.5–26.1	6–16	6	3	23	94	0.991
202 Croatia	$m^3$	cm	m	5–15	—	30	3		—	—
<i>Quercus laevis</i> (Turkey oak, Cer)										
203 Romania	$m^3$	cm	m	—	—	18	3		—	—
<i>Quercus pubescens</i> (Downy oak, Stejar pufos)										
204 Croatia	$ln(m^3)$	cm	m	4.1–23	3.5–15.1	2	3		347	0.96
205 Croatia	$ln(m^3)$	cm	m	4.1–23	3.5–15.1	2	3	23	347	0.932
206 Romania	$m^3$	cm	m	—	—	18	3		—	—
<i>Quercus robur</i> (Pedunculate oak)										
207 Netherlands	$dm^3$	cm	m	—	—	16	4		108	0.995
<i>Quercus rubra</i> (Red oak, Chêne rouge)										
208 Belgium	$m^3$	cm	m	—	—	14	6	2	891	—
209 Netherlands	$dm^3$	cm	m	—	—	16	4		793	0.996
<i>Quercus</i> spp. (Oak, Chênes, Stejar)										
210 Austria	$dm^3$	dm	dm	1—	—	29	5		—	—
211 Austria	$dm^3$	dm	dm	5–10.4	—	32	5		—	0.64
212 Belgium	$m^3$	cm	m	—	—	14	6	2	290	—
213 Netherlands	$dm^3$	mm	m	—	—	31	3	10	—	—
214 Netherlands	$dm^3$	mm	m	—	—	31	3	11	—	—
215 Netherlands	$dm^3$	mm	m	—	—	31	3	12	—	—
216 Romania	$m^3$	cm	m	—	—	18	3		—	—
217 UK	$m^3$	cm	m	—	—	8	2		—	—
<i>Salix caprea</i> (Goat willow, Salcie capreasca)										
218 Norway	$dm^3$	cm	m	5—	—	3	9		—	—
219 Romania	$m^3$	cm	m	—	—	18	3		—	—
<i>Salix</i> spp. (Salcie)										
220 Romania	$m^3$	cm	m	—	—	18	3		—	—
221 Romania	$m^3$	cm	m	—	—	18	3		—	—
<i>Sorbus aucuparia</i> (Rowan)										
222 Norway	$dm^3$	cm	m	5—	—	3	9		—	—
<i>Thuja plicata</i> (Western redcedar)										
223 Netherlands	$dm^3$	cm	m	—	—	16	4		165	0.993
224 Norway	$dm^3$	cm	m	5—	—	37	9	1	—	—
<i>Tilia cordata</i> (Tei)										
225 Romania	$m^3$	cm	m	—	—	18	3		—	—
<i>Tsuga heterophylla</i> (Hemlock)										
226 Netherlands	$dm^3$	cm	m	—	—	16	4		121	0.995
227 Norway	$dm^3$	cm	m	5—	—	37	9	1	—	—
<i>Ulmus</i> spp. (Elm, Orme, Ulm)										
228 Belgium	$m^3$	cm	m	—	—	14	6	2	276	—
229 Netherlands	$dm^3$	cm	m	—	—	16	4		108	0.996
230 Romania	$m^3$	cm	m	—	—	18	3		—	—

## Appendix B – References:

- 1 Bauger, E. 1995. Funksjoner og tabeller for kubering av stående trær. Rapport fra Skogforsk(16): 1–26.
- 2 Benko, M., Novotny, V., Vrbek, B. & Szirovicza, L. 2000. Volume tables of downy oak (*Quercus pubescens* Willd.). Radovi Sumarski Institut Jastrebarsko 8: 1–68.
- 3 Braastad, H. 1966. Volumtabeller for bjørk. Meddelelser fra det Norske Skogforsøksvesen 21(1): 23–78.
- 4 Brandel, G. 1974. Volymfunktioner för tall och gran. Skoghögskolan, Institutionen för skogsproduktion, Rapporter och Uppsatser 33: 178–191.
- 5 Brandel, G. 1990. Volumfunktioner för enskilda träd. Sveriges lantbruksuniversitet, Institutionen för skogsproduktion, Rapport 26: 1–181.
- 6 Brandini, P. & Tabacchi, G. 1996. Biomass and volume equations for holm oak and strawberry-tree in coppice stands of Southern Sardinia. ISAFA Communicazioni di Ricerca 96(1): 59–69.
- 7 Brantseg, A. 1967. Furu sønnafjells: kubering av stående skog, funksjoner og tabeller. Meddelelser fra det Norske Skogforsøksvesen 84: 689–739.
- 8 Broadmeadow, M. & Matthews, R. 2004. Survey methods for Kyoto Protocol monitoring and verification of UK forest carbon stocks. UK Emissions by Sources and Removals by Sinks due to Land Use, Land Use Change and Forestry Activities, Report (June 2004). CEH, Edinburgh.
- 9 Bruchwald, A. & Rymer-Dudzinska, T. 1987. Empirical equations for determining the volume of large timber of stem of growing spruce trees. Sylwan 131(5): 15–20.
- 10 Bruchwald, A. 1987. Empirical formulas for determining the stem volume of standing spruce trees. Annals of Warsaw Agricultural University – SGGW-AR Forestry and Wood Technology 36: 57–60.
- 11 Børset, O. 1954. Kubering av osp på rot. Meddelelser fra det norske Skogforsøksvesen 12: 391–447.
- 12 Cerný, M. 1990. Biomass of *Picea abies* (L.) Karst. in midwestern Bohemia. Scandinavian Journal of Forest Research 5: 83–95.
- 13 Corona, P. & Ferrara, A. 1987. Dendrometrical investigations on *Pinus silvestris* in Trentino-Alto Adige. Monti e Boschi 38(6): 51–54.
- 14 Dagnelie, P., Palm, R., Rondeux, J. & Thill, A. 1999. Tables de cubage des arbres et des peuplements forestiers. Les Presses Agronomiques de Gembloux, Gembloux. 126 p.
- 15 De Vries, P.G. 1961. The principle of nomograms applied to the stem volume functions of the volume tables for forest trees grown in the Netherlands. Nederlands Bosbouw Tijdschrift 33(5): 114–121.
- 16 Dik, E.J. 1984. Estimating the wood volume of standing trees in forestry practice. Rijksinstituut voor onderzoek in de bos en landschapsbouw de Dorschkamp, Wageningen. Uitvoerige verslagen 19(1): 1–114.
- 17 Eriksson, H. 1973. Volymfunktioner för stående träd av ask, asp, klibbal och contorta-tall. Institutionen för Skogsproduktion, Royal College of Forestry, Stockholm. Research Notes 26: 1–26.
- 18 Giurgiu, V. 1974. O expresie matematica unica a relatiei diametru – înaltime – volum, pentru majoritatea speciilor forestiere din Romania. Silvicultura si Exploatarea Padurilor 89(4): 173–178.
- 19 Hakkila, P. 1979. Wood density survey and dry weight tables for pine, spruce and birch stems in Finland. Communicationes Instituti Forestalis Fenniae 96(3): 1–59.
- 20 Hempel, G. 1968. Allometrische studie an *Pinus cembra* spp. *sibirica* (Rupr.) Kryl. und *Abies sibirica* (Ledeb.). Archiv für Forstwesen 17(11): 1099–1115.
- 21 Kanninen, K., Uusvaara, O. & Valonen, P. 1979. Kokopuuraaka-aineen mittaus ja ominaisuudet. Folia Forestalia 403: 1–53.
- 22 Laasasenaho, J. 1982. Taper curve and volume functions for pine, spruce and birch. Communicationes Instituti Forestalis Fenniae 108: 1–74.
- 23 Laasasenaho, J. & Sevola, Y. 1971. Mänty- ja kuusirunkojen puitavarasuhteet ja kantoarvot. Communicationes Instituti Forestalis Fenniae 74(3): 1–87.
- 24 Lockow, K.-W. 1993. Modellbildung und quantifizierung der durchmesser- und volumenstruktur des ausscheidenden kieferjungbestandes – holzmeßkundliche entscheidungshilfen für die erst-durchforstung. Beiträge für Forstwirtschaft und Landschaftsökologie 27(2): 77–82.
- 25 Mund, M., Kummetz, E., Hein, M., Bauer, G.A. & Schulze, E.-D. 2002. Growth and carbon stocks of a spruce forest chronosequence in central Europe. Forest Ecology and Management 171: 275–296.
- 26 Norrby, M. 1990. Volym- och formtalsfunktioner for *Larix sukaczewii* och *Larix sibirica* på Island.

- Institutionen för skogsskötsel. Series Volym- och formtalsfunktioner för Larix sukaczewii och Larix sibirica på Island. Sveriges Lantbruksuniversitet, Umeå.
- 27 Näslund, M. 1947. Funktioner och tabeller för kubering av stående träd. Meddelanden från Statens skogsforskningsinstitutet 36(3): 1–81.
- 28 Pellinen, P. 1986. Biomasseuntersuchungen im Kalkbuchenwald. Series Biomasseuntersuchungen im Kalkbuchenwald. University of Göttingen, Germany. 145 p.
- 29 Pollanschütz, J. 1974. Formzahlfunktionen der Hauptbaumarten Österreichs. Allgemeine Forstzeitung 85: 341–343.
- 30 Pranjic, A. & Lukic, N. 1986. Form factor and volume table for Holm oak (*Quercus ilex* L.). Glasnik za Sumske Pokuse, Posebno Izdanje 2: 169–177.
- 31 Schelhaas, M.J., Nabuurs, G.J., Jans, W.W.P., Moors, E.J., Sabaté, S. & Daamen, W.P. 2002. Converging estimates of the forest carbon sink. Alterra-rapport 631: 1–44.
- 32 Schieler, K. 1988. Diploma thesis, Institute for Forest growth and Yield Reserch, University for Agricultural Sciences, Vienna.
- 33 Snorrason, A. & Einarsson, S.F. 2004. Single-tree biomass and stem volume functions for eleven tree species used in Icelandic forestry. Submitted manuscript.
- 34 Vestjordet, E. 1967. Funksjoner og tabeller for kubering av stående gran. Meddelelser fra det Norske Skogforsksvesen 84: 539–574.
- 35 Vuokila, Y. 1965. Functions for variable density yield tables of pine based on temporary sample plots. Communicationes Instituti Forestalis Fenniae 60(4): 1–86.
- 36 Xiao, C.W., C.Y., J., Janssens, I.A., Roskams, P., Nachtergale, L., Carrara, A., Sanchez, B.Y. & Ceulemans, R. 2003. Above-and belowground biomass and net primary production in a 73-year-old Scots pine forest. Manuscript.
- 37 Øen, S., Bauger, E. & Øyen, B.-H. 2001. Functionar for volumberekning av framande treslag i Vest-Norge. Aktuelt fra Skogforsk 3/01: 18–19.

## Appendix B – Contact persons

- 1 **Joosten R.**, Joosten@t-online.de, Univertsty Paderborn, Germany
- 2 **Matthews, R.**, robert.matthews@forestry.gsi.gov.uk, Forest Research, Alice Holt Research Station, UK
- 3 **Muukkonen P.**, petteri.muukkonen@metla.fi, Finnish Forest Research Institute, Finland
- 4 **Nabuurs G.-J.**, G.J.Nabuurs@Alterra.wag-ur.nl, Alterra, Wageningen, the Netherlands
- 5 **Prskawetz M.**, michael.prskawetz@bfw.gv.at, Federal Office and Research Centre for Forests, Vienna, Austria
- 6 **Sheridan M.**, sheridan.m@fsagx.ac.be , Department of Plant Biology, Gembloux agricultural University, Belgium
- 7 **Snorrason A.**, arnor@skogur.is, Skógrækt ríkisins, Iceland
- 8 **Xiao C.W.**, Xiao@uia.ua.ac.be, University of Antwerpen, Belgium
- 9 **Zianis D.**, zianis.dimitris@ed.ac.uk, IERM, Edinburgh University, UK

## Appendix B – Comments

- 1 Plantations
- 2 Stem volume to upper girth limit of 22 cm
- 3 Including bark
- 4 Excluding bark
- 5 Merchantable volume >34 cm
- 6 Including bark, Southern Sweden, adjusted  $r^2$
- 7 Excluding bark, Southern Sweden, adjusted  $r^2$
- 8 Including bark, Northern Sweden, adjusted  $r^2$
- 9 Excluding bark, Northern Sweden, adjusted  $r^2$
- 10 Dominant
- 11 Dominated
- 12 Seed trees
- 13 Merchantable wood >7 cm
- 14 Non merchantable wood <7 cm
- 15 Including bark, trees without forks
- 16 Including bark, forked trees
- 17 Excluding bark, trees without forks
- 18 Excluding bark, forked trees
- 19 Without bark
- 20 Tree height 4 m and more
- 21 Tree height over 10 m
- 22 Merchantable volume
- 23 Merchantable volume, >3–4 cm

**Appendix C.** Volume equations for different tree species. The format of the stem volume equation is given in the column labelled Equation, and a–g are parameter values. The “ln” is the natural logarithm and the “log” is the 10-based logarithm.

	Equation
<i>Abies alba</i> (Silver fir)	
1 Norway	$a \cdot H^b \cdot D^c \cdot (H-1.3)^d \cdot (D+100)^e$
<i>Abies grandis</i> (Grand fir)	
2 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
3 Norway	$a \cdot H^b \cdot D^c \cdot (H-1.3)^d \cdot (D+100)^e$
<i>Abies sibirica</i> (Fir)	
4 Germany	$a \cdot D^b$
<i>Abies</i> spp. (Fir, Brad)	
5 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 \cdot H \cdot \ln(D)^2 + c \cdot D^2 + d \cdot D \cdot H + e \cdot H + f \cdot D + g)$
6 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 \cdot H \cdot \ln(D)^2 + c \cdot D^2 + d \cdot D \cdot H)$
7 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
<i>Acacia</i> spp. (Salcim)	
8 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
<i>Acer pseudoplatanus</i> (Maple, Erable sycomore, Paltin, Sycamore)	
9 Belgium	$a+b \cdot D+c \cdot D^2+d \cdot D^3+e \cdot H+f \cdot D^2 \cdot H$
10 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
11 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
12 UK	$a+b \cdot D^2 \cdot H^c$
<i>Alnus alba</i> (Anin alb)	
13 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
<i>Alnus glutinosa</i> (Black alder, Klibbal)	
14 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
15 Norway	$a+b \cdot D^2+c \cdot D^2 \cdot H$
16 Norway	$a+b \cdot D^2+c \cdot D^2 \cdot H+d \cdot H^2 \cdot D$
17 Sweden	$a \cdot D^2+b \cdot D^2 \cdot H+c \cdot D \cdot H^2+d \cdot D \cdot H+e \cdot D^2 \cdot H^2$
18 Sweden	$a \cdot D^b \cdot H^c$
19 Sweden	$a \cdot D^2+b \cdot D^2 \cdot H+c \cdot D \cdot H^2+d \cdot D \cdot H+e \cdot D^2 \cdot H^2$
<i>Alnus incana</i> (Grey alder)	
20 Norway	$a+b \cdot D^2+c \cdot D^2 \cdot H+d \cdot H^2 \cdot D+e \cdot H^2$
<i>Alnus nigra</i> (Anin negru)	
21 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
<i>Alnus</i> spp. (Alder)	
22 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 + c \cdot D)$
<i>Arbutus unedo</i> (Strawberry-tree)	
23 Italy	$a+b \cdot D^2$
<i>Betula pendula</i> (Birch, Berk)	
24 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
<i>Betula</i> spp. (Birch, Björk, Bjørk, Bouleaux, Mesteacan)	
25 Belgium	$a+b \cdot D+c \cdot D^2+d \cdot D^3+e \cdot H+f \cdot D^2 \cdot H$
26 Finland	$a+b \cdot \ln(D)$
27 Finland	$a+b \cdot \ln(c+d \cdot D)+e \cdot D$
28 Finland	$a \cdot (D^b) \cdot (c^D) \cdot (H^d) \cdot (H-1.3)^e$
29 Finland	$a+b \cdot \ln(D)+c \cdot \ln(H)+d \cdot \ln(H-1.3)+e \cdot D$
30 Norway	$a+b \cdot D^2+c \cdot D^2 \cdot H+d \cdot D \cdot H^2+e \cdot H^2$
31 Norway	$a+b \cdot D^2+c \cdot D^2 \cdot H+d \cdot D \cdot H^2+e \cdot H^2$
32 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
33 Sweden	$a \cdot D^2+b \cdot D^2 \cdot H+c \cdot D \cdot H^2+d \cdot H^2$
34 Sweden	$10^a \cdot D^b \cdot (D+20)^c \cdot H^d \cdot (H-1.3)^e$
35 Sweden	$10^a \cdot D^b \cdot (D+20)^c \cdot H^d \cdot (H-1.3)^e$
36 Sweden	$10^a \cdot D^b \cdot (D+20)^c \cdot H^d \cdot (H-1.3)^e$

	a	b	c	Parameters d	e	f	g
1	1.6662	3.2394	1.9334	-1.8997	-0.9739	-	-
2	1.77220	0.96736	-2.45224	-	-	-	-
3	1.6662	3.2394	1.9334	-1.8997	-0.9739	-	-
4	0.0001316	2.52	-	-	-	-	-
5	0.580223	-0.0307373	-17.1507	0.089869	-0.080557	19.661	-2.4584
6	0.560673	0.15468	-0.65583	0.033210	-	-	-
7	4.52·10 <sup>-5</sup>	2.1554	-0.1067	0.9380	0.0228	-	-
8	0.00046903	1.807	0.0292	-0.4155	0.5455	-	-
9	0.010343	-0.00450536	3.4070·10 <sup>-4</sup>	-4.0472·10 <sup>-6</sup>	7.7115·10 <sup>-4</sup>	2.9836·10 <sup>-5</sup>	-
10	1.89756	0.97716	-2.94253	-	-	-	-
11	0.00035375	1.02	0.3997	0.666	0.021	-	-
12	-0.012668	0.0000737	0.75	-	-	-	-
13	0.00065013	1.6750	0.1001	-0.4990	0.5902	-	-
14	1.85749	0.88675	-2.5222	-	-	-	-
15	8.6524	0.076844	0.031573	-	-	-	-
16	0.6716	0.75708	0.029679	0.004341	-	-	-
17	0.1926	0.01631	0.003755	-0.02756	0.000499	-	-
18	0.05437	1.94505	0.92947	-	-	-	-
19	0.2264	0.01347	0.007665	-0.06669	0.000428	-	-
20	-1.86827	0.21461	0.01283	0.0138	-0.06311	-	-
21	8.666·10 <sup>-5</sup>	1.7148	0.1014	0.801	0.0530	-	-
22	0.387399	7.17123	0.04407	-	-	-	-
23	-0.5547	0.3757	-	-	-	-	-
24	1.89060	.26595	-1.07055	-	-	-	-
25	-0.011392	-0.00031447	0.000279211	-5.7966·10 <sup>-6</sup>	-5.9573·10 <sup>-4</sup>	3.0409·10 <sup>-5</sup>	-
26	-2.09787	2.55058	-	-	-	-	-
27	-5.41948	3.57630	2	1.25	-0.0395855	-	-
28	0.011197	2.10253	0.986	3.98519	-2.65900	-	-
29	-4.4759	2.0851	4.0691	-2.7375	-0.013311	-	-
30	-1.86827	0.21461	0.01283	0.01380	0.06311	-	-
31	0.99983	0.006325	0.02849	0.00885	-0.00799	-	-
32	8.141·10 <sup>-5</sup>	2.248	-0.2062	0.1946	0.4147	-	-
33	0.1305	0.01338	0.01757	-0.05606	-	-	-
34	-0.89359	2.27954	-1.18672	7.07362	-5.45175	-	-
35	-0.93631	2.30212	-1.40378	8.01817	-6.18825	-	-
36	-0.44224	2.47580	-1.40854	5.16863	-3.77147	-	-

## App. C

	Equation
37 Sweden	$10^a \cdot D^b \cdot (D+20)^c \cdot H^d \cdot (H-1.3)^e$
38 Sweden	$a \cdot D^2 + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2$
39 UK	$a+b \cdot D^2 \cdot H^c$
<i>Carpinus</i> spp.	
40 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
41 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
42 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
<i>Chamaecyparis lawsoniana</i> (Lawson cypress)	
43 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
<i>Corylus avellana</i> (Hazel)	
44 Norway	$a+b \cdot D^2 + c \cdot D^2 \cdot H + d \cdot D \cdot H^2 + e \cdot H^2$
<i>Fagus</i> spp. (Beech, Fag)	
45 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 \cdot H \cdot \ln(D)^2 + c \cdot D^2 + d \cdot D \cdot H + e \cdot H + f \cdot D + g)$
46 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 + c \cdot D)$
47 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D))^2 + d \cdot \log(H) + e \cdot \log(H)^2}$
48 UK	$a+b \cdot D^2 \cdot H^c$
<i>Fagus sylvatica</i> (Beech, Rotbuche, Beuk)	
49 Belgium	$a+b \cdot D+c \cdot D^2+d \cdot D^3+e \cdot H+f \cdot D^2 \cdot H$
50 Germany	$a+b \cdot D \cdot H^2+c \cdot D^3$
51 Germany	$a+b \cdot D \cdot H^2+c \cdot D^3$
52 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
53 Netherlands	$a \cdot D^b \cdot H^c$
<i>Fraxinus excelsior</i> (Ash, Frêne, Es)	
54 Belgium	$a+b \cdot D+c \cdot D^2+d \cdot D^3+e \cdot H+f \cdot D^2 \cdot H$
55 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
56 Sweden	$a \cdot D^2 \cdot H+b \cdot D^2+c \cdot D \cdot H$
57 Sweden	$a \cdot D^2 \cdot H+b \cdot D^2+c \cdot D \cdot H$
58 Sweden	$a \cdot D^b \cdot H^c$
59 Sweden	$a \cdot D^2 \cdot H+b \cdot D^2+c \cdot D \cdot H$
60 Sweden	$a \cdot D^2 \cdot H+b \cdot D^2+c \cdot D \cdot H$
<i>Fraxinus</i> spp. (Ash)	
61 Norway	$a+b \cdot D^2+c \cdot D^2 \cdot H+d \cdot D \cdot H^2+e \cdot H^2$
62 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D))^2 + d \cdot \log(H) + e \cdot \log(H)^2}$
63 UK	$a+b \cdot D^2 \cdot H^c$
<i>Larix decidua</i> (Larch, Mélèzes)	
64 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 \cdot H \cdot \ln(D)^2 + c \cdot D^2 + d \cdot D \cdot H + e \cdot H + f \cdot D + g)$
65 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 + c \cdot D)$
66 Belgium	$a+b \cdot D+c \cdot D^2+d \cdot D^3+e \cdot H+f \cdot D^2 \cdot H$
67 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
68 Norway	$a \cdot H^b \cdot D^c \cdot (H-1.3)^d \cdot (D+100)^e$
<i>Larix hybrid</i> (Hybrid larch)	
69 Norway	$a \cdot H^b \cdot D^c \cdot (H-1.3)^d \cdot (D+100)^e$
<i>Larix kaempferi</i> (Japanese larch)	
70 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
71 Norway	$a \cdot H^b \cdot D^c \cdot (H-1.3)^d \cdot (D+100)^e$
<i>Larix sibirica</i> (Siberian larch)	
72 Iceland	$\exp(a) \cdot D^b \cdot H^c$
73 Iceland	$\exp(a) \cdot D^b \cdot H^c$
74 Iceland	$a \cdot D^b \cdot H^c$
75 Norway	$a \cdot H^b \cdot D^c \cdot (H-1.3)^d \cdot (D+100)^e$
<i>Larix</i> spp. (Lehtikuusi, Lork, Larice)	
76 Finland	$a+b \cdot D^2+c \cdot D^2 \cdot H+d \cdot H^2 \cdot D$
77 Finland	$a+b \cdot D^2+c \cdot D^2 \cdot H+d \cdot H^2 \cdot D$

**App. C**

	a	b	c	Parameters d	e	f	g
37	-0.35394	2.52141	-1.54257	4.88165	-3.47422	-	-
38	0.1432	0.008561	0.02180	-0.06630	-	-	-
39	-0.009184	0.0000673	0.75	-	-	-	-
40	0.00021491	2.258957614	0.001411006	0.60291075	-	-	-
41	0.00021491	2.258957614	-0.01120638	0.60291075	-	-	-
42	0.00021491	2.258957614	-0.00956695	0.60291075	-	-	-
43	1.85298	.86717	-2.33706	-	-	-	-
44	-1.86827	0.21461	0.01283	0.0138	-0.06311	-	-
45	0.989253	-0.0371508	-31.0674	-0.386321	0.219462	49.6136	-22.372
46	0.517300	-13.62144	9.9888	-	-	-	-
47	0.0000757	1.3791	0.2127	1.1992	-0.0584	-	-
48	-0.014306	0.0000748	0.75	-	-	-	-
49	-0.015572	0.00290013	$-7.0476 \cdot 10^{-6}$	$2.393 \cdot 10^{-6}$	-0.0013528	$3.9837 \cdot 10^{-5}$	-
50	$15.589 \cdot 10^{-3}$	$0.01696 \cdot 10^{-3}$	$0.01883 \cdot 10^{-3}$	-	-	-	-
51	$16.641 \cdot 10^{-3}$	$0.72179 \cdot 10^{-3}$	$0.00252 \cdot 10^{-3}$	-	-	-	-
52	1.55448	1.55880	-3.57875	-	-	-	-
53	0.049	1.78189	1.08345	-	-	-	-
54	-0.039836	0.006262765	-0.00015937	$-1.9902 \cdot 10^{-7}$	-0.0009834	$3.7872 \cdot 10^{-5}$	-
55	1.95277	0.77206	-2.48079	-	-	-	-
56	0.03246	0.03310	0.04127	-	-	-	-
57	0.03593	0.03310	0.04127	-	-	-	-
58	0.06328	1.92428	0.88690	-	-	-	-
59	0.03249	0.02941	0.03892	-	-	-	-
60	0.03453	0.02941	0.03892	-	-	-	-
61	-1.86827	0.21461	0.01283	0.0138	-0.06311	-	-
62	0.00030648	1.2676	0.3102	0.4929	0.0962	-	-
63	-0.012107	0.0000777	0.75	-	-	-	-
64	0.609443	-0.0455748	-18.6631	-0.248736	0.126594	36.9783	-14.204
65	0.487270	-2.04291	5.9995	-	-	-	-
66	-0.03088	0.004676261	$-4.8614 \cdot 10^{-5}$	$-3.8178 \cdot 10^{-6}$	-0.0011638	$4.0597 \cdot 10^{-5}$	-
67	1.86670	1.08118	-3.0488	-	-	-	-
68	0.7761	3.6461	1.9166	-2.3179	-0.8236	-	-
69	0.7761	3.6461	1.9166	-2.3179	-0.8236	-	-
70	1.87077	1.00616	-2.8748	-	-	-	-
71	0.7606	3.5377	1.9741	-2.1902	-0.8459	-	-
72	-2.5079	1.7574	0.9808	-	-	-	-
73	-2.9946	1.8105	0.9908	-	-	-	-
74	0.0983	1.551	1.1483	-	-	-	-
75	0.7761	3.6461	1.9166	-2.3179	-0.8236	-	-
76	0.5	0.0753	0.03345	-0.00243	-	-	-
77	0.4	-0.01	0.03355	-0.00359	-	-	-

## App. C

	Equation
78 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
79 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
80 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
81 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
<i>Picea abies</i> (Norway spruce, Kuusi, Gran, Epicéa, Fijnspar)	
82 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 \cdot H \cdot \ln(D)^2 + c \cdot D^2 + d \cdot D \cdot H + e \cdot H + f \cdot D)$
83 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 \cdot H \cdot \ln(D)^2 + c \cdot D^2 + d \cdot D)$
84 Belgium	$a + b \cdot D + c \cdot D^2 + d \cdot D^3 + e \cdot H + f \cdot D^2 \cdot H$
85 Czech Republic	$a \cdot D^b$
86 Czech Republic	$a \cdot (H \cdot D^2)^b$
87 Finland	$a + b \cdot \ln(D)$
88 Finland	$a + b \cdot \ln(c + d \cdot D) + e \cdot D$
89 Finland	$a \cdot (D^b) \cdot (c^D) \cdot (H^d) \cdot (H - 1.3)^e$
90 Finland	$a + b \cdot \ln(D) + c \cdot \ln(H) + d \cdot \ln(H - 1.3) + e \cdot D$
91 Finland	$a \cdot D^b \cdot H^c$
92 Finland	$a \cdot D^b \cdot H^c$
93 Finland	$a + b \cdot \ln(D) + c \cdot D^2$
94 Germany	$a \cdot H \cdot D^2$
95 Germany	$a \cdot H \cdot D^2$
96 Iceland	$a \cdot D^b \cdot H^c$
97 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
98 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
99 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
100 Netherlands	$D^a \cdot H^b \cdot e^c$
101 Netherlands	$a \cdot D^b \cdot H^c$
102 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 40)^e$
103 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 40)^e$
104 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 40)^e$
105 Norway	$a + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2 + e \cdot D \cdot H$
106 Norway	$a + b \cdot D \cdot H^2 + c \cdot H^2 + d \cdot D \cdot H + e \cdot H + f \cdot D$
107 Norway	$a + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2 + e \cdot D \cdot H$
108 Norway	$a + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2 + e \cdot D \cdot H$
109 Norway	$a + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2 + e \cdot D \cdot H$
110 Norway	$a + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot D^2$
111 Norway	$a + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2 + e \cdot D \cdot H$
112 Norway	$a + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2 + e \cdot D \cdot H$
113 Norway	$a + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2 + e \cdot D \cdot H$
114 Poland	$(\pi/40000) \cdot H \cdot D \cdot (a + b \cdot D)$
115 Poland	$(\pi/40000) \cdot H \cdot D \cdot (a \cdot D + b)$
116 Sweden	$a \cdot D^2 + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2$
117 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$
118 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$
119 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$
120 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$
121 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$
122 Sweden	$a \cdot D^2 + b \cdot D^2 \cdot H + c \cdot D \cdot H^2 + d \cdot H^2$
123 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$
<i>Picea engelmannii</i> (Engelmanni spruce)	
124 Iceland	$a \cdot D^b \cdot H^c$
<i>Picea sitchensis</i> (Sitka spruce)	
125 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
126 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 40)^e$
127 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 40)^e$

## App. C

	a	b	c	Parameters d	e	f	g
78	0.00035217	2.12841828	0.003292718	0.76283925	—	—	—
79	0.00035217	2.12841828	-0.1054168	0.76283925	—	—	—
80	0.00035217	2.12841828	-0.0026067	0.76283925	—	—	—
81	2.822·10 <sup>-5</sup>	2.2060	-0.1136	1.115	0.0129	—	—
82	0.46818	-0.013919	-28.213	0.37474	-0.28875	28.279	—
83	0.563443	-0.12731	-8.55022	7.6331	—	—	—
84	-0.010929	0.004380951	-9.4713·10 <sup>-5</sup>	-7.8024·10 <sup>-6</sup>	-0.0027922	4.8346·10 <sup>-5</sup>	—
85	0.00059707	2.1286	—	—	—	—	—
86	0.00011261	0.87852	—	—	—	—	—
87	-2.41218	2.62463	—	—	—	—	—
88	-5.39934	3.46468	2	1.25	-0.0273199	—	—
89	0.022927	1.91505	0.99146	2.82541	-1.53547	—	—
90	-3.7544	1.8960	2.8979	-1.6020	-0.007827	—	—
91	0.7877	1.9302	0.79465	—	—	—	—
92	0.10838	1.8202	0.77154	—	—	—	—
93	-2.59385	2.71757	-0.000097	—	—	—	—
94	0.502	—	—	—	—	—	—
95	0.418	—	—	—	—	—	—
96	0.1299	1.6834	0.8598	—	—	—	—
97	0.00053238	2.164126647	0.004108377	0.54879808	—	—	—
98	0.00053238	2.164126647	-0.04670018	0.54879808	—	—	—
99	0.00053238	2.164126647	-0.0102582	0.54879808	—	—	—
100	1.75055	1.10897	-2.75863	—	—	—	—
101	0.04143	1.6704	1.3337	—	—	—	—
102	0.6844	3.0296	2.0560	-1.7377	-0.9756	—	—
103	0.7464	2.496	2.0714	-1.4175	-0.9601	—	—
104	0.5824	1.1987	1.9339	-0.0594	-0.7442	—	—
105	0.52	0.02403	0.01463	-0.10983	0.15195	—	—
106	-31.57	0.0016	0.0186	0.63	-2.34	3.2	—
107	10.14	0.0124	0.03117	-0.36381	0.28578	—	—
108	6.69	0.01308	0.02853	-0.31956	0.28969	—	—
109	0.46	0.02427	0.01521	-0.18254	0.20994	—	—
110	0.67	0.03023	0.00712	0.04175	—	—	—
111	0.28	0.00815	0.03053	-0.50725	0.51643	—	—
112	0.3	0.02593	0.01268	-0.0977	0.14586	—	—
113	4.33	0.01491	0.02606	-0.31854	0.31106	—	—
114	0.666151	0.458507	—	—	—	—	—
115	0.53005	1.229283	—	—	—	—	—
116	0.1150	0.01746	0.02022	-0.05618	—	—	—
117	-0.9513	1.9781	-0.5254	2.7604	-1.4684	—	—
118	-0.79783	2.07157	-0.73882	3.16332	-1.82622	—	—
119	-1.02039	2.00128	-0.47473	2.87138	-1.61803	—	—
120	-0.82249	2.11094	-0.89626	3.51812	-2.05567	—	—
121	-1.06019	2.04239	-0.54292	2.80843	-1.52110	—	—
122	0.1104	0.01925	0.01815	-0.04936	—	—	—
123	-1.0342	1.9683	-0.3850	2.4018	-1.2075	—	—
124	0.4693	1.311	0.781	—	—	—	—
125	1.78383	1.13397	-2.90893	—	—	—	—
126	0.1614	3.7060	1.9747	-2.2905	-0.6665	—	—
127	0.1870	3.7077	1.9854	-2.2816	-0.7161	—	—

## App. C

	Equation
128 Norway	$a \cdot D^b \cdot (H - 1.3)^c \cdot (D + 40)^d$
<i>Picea</i> spp. (Molid)	
129 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
130 Iceland	$a \cdot D^b \cdot H^c$
<i>Pinus contorta</i> (Contorta tall)	
131 Iceland	$a \cdot D^b \cdot H^c$
132 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
133 Sweden	$a \cdot D^2 + b \cdot D^2 H + c \cdot D^2 H^2 - d \cdot D \cdot H + e \cdot D \cdot H^2$
134 Sweden	$a \cdot D^b \cdot H^c$
135 Sweden	$a \cdot D^2 + b \cdot D^2 H + c \cdot D^2 H^2 - d \cdot D \cdot H + e \cdot D \cdot H^2$
<i>Pinus nigra</i> var <i>maritima</i> (Black pine)	
136 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
<i>Pinus nigra</i> var <i>nigra</i> (Black pine, Pin negru)	
137 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
138 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
<i>Pinus</i> spp.	
139 Germany	$a \cdot H^b$
140 Germany	$a \cdot D^b$
141 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
142 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
143 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
<i>Pinus sylvestris</i> (Scots pine, Mänty, Tall, Furu, Grove den, Pin silvestri)	
144 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 \cdot H \cdot \ln(D)^2 + c \cdot D^2 + d \cdot H)$
145 Belgium	$a \cdot D^b$
146 Belgium	$a + b \cdot D + c \cdot D^2 + d \cdot D^3 + e \cdot H + f \cdot D^2 \cdot H$
147 Finland	$a + b \cdot \ln(D)$
148 Finland	$a + b \cdot \ln(c + d \cdot D) + e \cdot D$
149 Finland	$a \cdot (D^b) \cdot (D^d) \cdot (H - 1.3)^e$
150 Finland	$a + b \cdot \ln(D) + c \cdot \ln(H) + d \cdot \ln(H - 1.3) + e \cdot D$
151 Finland	$a \cdot D^b \cdot H^c$
152 Finland	$a \cdot D^b \cdot H^c$
153 Finland	$a \cdot H \cdot D^2 + b \cdot D \cdot H + c \cdot D^3 + d \cdot D \cdot H^2$
154 Finland	$a + b \cdot \ln(D) + c \cdot D^2$
155 Germany	$a \cdot D^b \cdot H^c$
156 Italy	$a \cdot D^b \cdot H^c$
157 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
158 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
159 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
160 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
161 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 100)^e$
162 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 100)^e$
163 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 100)^e$
164 Norway	$a + b \cdot D^2 + c \cdot D^2 \cdot H$
165 Norway	$a + b \cdot D^2 + c \cdot D^2 \cdot H + d \cdot D \cdot H^2$
166 Norway	$a + b \cdot D^2 + c \cdot D^2 \cdot H$
167 Norway	$a + b \cdot D^2 + c \cdot D^2 \cdot H$
168 Norway	$a + b \cdot D^2 \cdot H$
169 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
170 Sweden	$a \cdot D^2 + b \cdot D^2 \cdot H + c \cdot D \cdot H^2$
171 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$
172 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$
173 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$
174 Sweden	$10^a \cdot D^b \cdot (D + 20)^c \cdot H^d \cdot (H - 1.3)^e$

## App. C

	a	b	c	Parameters d	e	f	g
128	0.2101	1.8920	1.1095	-0.3895	-	-	-
129	0.00009464	1.9341	-0.0722	0.6365	0.172	-	-
130	0.0739	1.7508	1.0228	-	-	-	-
131	0.1491	1.6466	0.8325	-	-	-	-
132	1.89303	0.98667	-2.88614	-	-	-	-
133	0.1121	0.02870	-0.000061	0.09176	0.01249	-	-
134	0.04514	1.9005	1.06964	-	-	-	-
135	0.0883	0.03202	-0.000114	-0.07892	0.01049	-	-
136	1.89192	0.95374	-2.72505	-	-	-	-
137	1.95645	0.88671	-2.7675	-	-	-	-
138	0.00010892	1.9701	0.0102	0.4858	0.1330	-	-
139	0.000074	3.1	-	-	-	-	-
140	0.0001078	2.56	-	-	-	-	-
141	0.00042613	2.066225947	-0.001926657	0.80636901	-	-	-
142	0.00042613	2.066225947	-0.07956244	0.80636901	-	-	-
143	0.00042613	2.066225947	0.00369501	0.80636901	-	-	-
144	0.435949	-0.0149083	5.21091	0.028702	-	-	-
145	0.000244	2.32716	-	-	-	-	-
146	-0.039836	$4.8710 \cdot 10^{-3}$	$-6.1028 \cdot 10^{-5}$	$1.4889 \cdot 10^{-5}$	$7.3997 \cdot 10^{-5}$	$2.9221 \cdot 10^{-5}$	-
147	-2.2945	2.57025	-	-	-	-	-
148	-5.39417	3.48060	2	1.25	-0.039884	-	-
149	0.036089	2.01395	0.99676	2.07025	-1.07209	-	-
150	-3.2890	1.9995	2.1395	-1.1411	-0.002847	-	-
151	0.0942	1.9671	0.7005	-	-	-	-
152	0.095	1.9185	0.7381	-	-	-	-
153	0.05782	0.11632	-0.01092	-0.01317	-	-	-
154	-2.37912	2.62903	-0.000126	-	-	-	-
155	$5.6537 \cdot 10^{-5}$	1.960466	0.894433	-	-	-	-
156	1.480589	1.982459514	0.742674501	-	-	-	-
157	0.00207765	1.952764402	$-8.6651 \cdot 10^{-5}$	0.48560878	-	-	-
158	0.00207765	1.952764402	-0.11110535	0.48560878	-	-	-
159	0.00207765	1.952764402	0.001095496	0.48560878	-	-	-
160	1.82075	1.07427	-2.8885	-	-	-	-
161	0.1424	2.0786	1.9028	-1.0259	-0.2640	-	-
162	0.1263	2.4621	1.9008	-1.3716	-0.2663	-	-
163	0.4434	4.9667	1.9912	-3.6612	-0.7502	-	-
164	8.6524	0.076844	0.031573	-	-	-	-
165	0.6716	0.075708	0.029679	-	-	-	-
166	2.0044	0.029886	0.036972	0.004341	-	-	-
167	2.9121	0.039994	-0.001091	-	-	-	-
168	2.9361	0.038906	-	-	-	-	-
169	0.00014808	1.8341	-0.0448	0.3115	0.3525	-	-
170	0.1028	0.02705	0.005215	-	-	-	-
171	-1.1226	2.0180	-0.2135	1.8271	-0.8297	-	-
172	-1.20914	1.94740	-0.05947	1.40958	-0.45810	-	-
173	-1.38903	1.84493	0.06563	2.02122	-1.01095	-	-
174	-1.25246	1.98244	-0.13118	1.03781	-0.03482	-	-

## App. C

	Equation
175 Sweden	$10^a \cdot D^b \cdot (D+20)^c \cdot H^d \cdot (H-1.3)^e$
176 Sweden	$a \cdot D^2 + b \cdot D^2 \cdot H + c \cdot D \cdot H^2$
177 Sweden	$10^a \cdot D^b \cdot (D+20)^c \cdot H^d \cdot (H-1.3)^e$
<i>Populus</i> spp. (Poplar, Plop)	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 + c \cdot D)$
178 Austria	$a \cdot D^{(b+c)} \cdot H^d$
179 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
180 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
181 Netherlands	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
182 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
183 Romania	$a \cdot b \cdot D^2 \cdot H^c$
184 UK	
<i>Populus tremula</i> (Aspen, Plop tremulator)	$a + b \cdot D^2 \cdot H$
185 Norway	$a + b \cdot D^2 \cdot H$
186 Norway	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
187 Romania	$a \cdot D^2 + b \cdot D^2 \cdot H + c \cdot D^2 \cdot H^2 + d \cdot D \cdot H + e \cdot D \cdot H^2$
188 Sweden	$a \cdot D^b \cdot H^c$
189 Sweden	$a \cdot D^2 \cdot H + b \cdot D^2 \cdot H + c \cdot D^2 \cdot H^2 + d \cdot D \cdot H + e \cdot D \cdot H^2$
190 Sweden	
<i>Populus trichocarpa</i> (Black cottonwood)	$a \cdot D^b \cdot H^c$
191 Iceland	
<i>Prunus avium</i> (Wild cherry, Merisier, Zoete kers)	$a + b \cdot D + c \cdot D^2 + d \cdot D^3 + e \cdot H + f \cdot D^2 \cdot H$
192 Belgium	
<i>Pseudotsuga menziesii</i> (Douglas fir, Duglas)	$a + b \cdot D + c \cdot D^2 + d \cdot D^3 + e \cdot H + f \cdot D^2 \cdot H$
193 Belgium	$D^a \cdot H^b \cdot \exp(c)$
194 Netherlands	$a \cdot H^b \cdot D^c \cdot (H-1.3)^d \cdot (D+40)^e$
195 Norway	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
196 Romania	
<i>Pseudotsuga</i> spp.	$a \cdot D^{(b+c)} \cdot H^d$
197 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
198 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
199 Netherlands	
<i>Quercus grisea</i> (Gray oak, Stejar brumariu)	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
200 Romania	
<i>Quercus ilex</i> (Holm oak)	$a + b \cdot D^2 \cdot H$
201 Italy	$a \cdot D^b \cdot H^c$
202 Croatia	
<i>Quercus laevis</i> (Turkey oak, Cer)	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
203 Romania	
<i>Quercus pubescens</i> (Downy oak, Stejar pufos)	$a + b \cdot \ln(D) + c \cdot H$
204 Croatia	$a + b \cdot \ln(D) + c \cdot H$
205 Croatia	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
206 Romania	
<i>Quercus robur</i> (Pedunculate oak)	$D^a \cdot H^b \cdot \exp(c)$
207 Netherlands	
<i>Quercus rubra</i> (Red oak, Chêne rouge)	$a + b \cdot D + c \cdot D^2 + d \cdot D^3 + e \cdot H + f \cdot D^2 \cdot H$
208 Belgium	$D^a \cdot H^b \cdot \exp(c)$
209 Netherlands	
<i>Quercus</i> spp. (Oak, Chênes, Stejar)	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 + c \cdot D \cdot H + d \cdot H + e \cdot D + f)$
210 Austria	$(\pi/4) \cdot (a \cdot D^2 \cdot H + b \cdot D^2 \cdot H \cdot \ln(D)^2 + c \cdot D^2)$
211 Austria	$a + b \cdot D + c \cdot D^2 + d \cdot D^3 + e \cdot H + f \cdot D^2 \cdot H$
212 Belgium	$a \cdot D^{(b+c)} \cdot H^d$
213 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
214 Netherlands	

## App. C

	a	b	c	Parameters d	e	f	g
175	-1.52761	1.82928	0.07454	1.43792	-0.35559	-	-
176	0.1072	0.02427	0.007315	-	-	-	-
177	-1.2605	1.9322	-0.0897	2.1795	-1.1676	-	-
178	0.366419	1.13323	0.1306	-	-	-	-
179	0.0009507	1.895629295	0.001650837	0.8392146	-	-	-
180	0.0009507	1.895629295	-0.09208823	0.8392146	-	-	-
181	0.0009507	1.895629295	-0.00773694	0.8392146	-	-	-
182	0.00018059	1.9342	0.0013	-0.0161	0.4099	-	-
183	0.00041486	1.4466	0.1089	-0.1963	0.5681	-	-
184	-0.004298	0.0000435	0.89	-	-	-	-
185	9.69	0.0365	-	-	-	-	-
186	-0.21	0.0398	-	-	-	-	-
187	0.00007604	1.7812	0.0528	0.8533	0.0654	-	-
188	0.01548	0.03255	-0.000047	-0.01333	0.004859	-	-
189	0.03597	1.84297	1.15988	-	-	-	-
190	0.03392	-0.01491	-0.000005	0.01704	0.002926	-	-
191	0.0732	1.6933	1.0562	-	-	-	-
192	-0.002311	-0.00117728	0.000149061	$-7.8058 \cdot 10^{-6}$	$3.3282 \cdot 10^{-4}$	$3.1526 \cdot 10^{-5}$	-
193	-0.019911	0.001871101	0.000127328	$-5.7631 \cdot 10^{-6}$	0.00071591	$3.9371 \cdot 10^{-5}$	-
194	1.90053	.80726	-2.43151	-	-	-	-
195	1.8211	4.153	2.1342	-2.6902	-1.4265	-	-
196	$4.477 \cdot 10^{-5}$	1.8688	0.0424	1.1411	-0.1047	-	-
197	0.00095916	2.092560524	0.000297255	0.48824344	-	-	-
198	0.00095916	2.092560524	-0.0449007	0.48824344	-	-	-
199	0.00095916	2.092560524	0	0.48824344	-	-	-
200	$7.188 \cdot 10^{-5}$	1.4486	0.0204	1.4084	0.0409	-	-
201	1.1909	0.038639	-	-	-	-	-
202	0.000096	1.821	0.759	-	-	-	-
203	0.0001992	2.014	-0.0602	-0.1108	0.4811	-	-
204	-9.646	2.076	0.761	-	-	-	-
205	-11.473	2.548	0.967	-	-	-	-
206	0.00035164	1.1119	0.3108	0.5356	0.2139	-	-
207	2.00333	0.85925	-2.86353	-	-	-	-
208	-0.02149	0.002986681	$-4.2506 \cdot 10^{-5}$	$-2.1806 \cdot 10^{-6}$	-0.000743	$3.7473 \cdot 10^{-5}$	-
209	1.83932	0.9724	-2.71877	-	-	-	-
210	0.115631	65.9961	1.20321	-0.930406	-215.758	168.477	-
211	0.417118	0.21941	13.32594	-	-	-	-
212	-0.0022735	0.000389557	0.000124772	$-1.8434 \cdot 10^{-6}$	-0.0016657	$3.6985 \cdot 10^{-5}$	-
213	0.00095853	2.040672356	0.001965013	0.56366437	-	-	-
214	0.00095853	2.040672356	-0.02101921	0.56366437	-	-	-

## App. C

	Equation
215 Netherlands	$a \cdot D^{(b+c)} \cdot H^d$
216 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
217 UK	$a + b \cdot D^2 \cdot H^c$
<i>Salix caprea</i> (Goat willow, Salcie capreasca)	
218 Norway	$a + b \cdot D^2 + c \cdot D^2 \cdot H + d \cdot H^2 \cdot D + e \cdot H^2$
219 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
<i>Salix</i> spp. (Salcie)	
220 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
221 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
<i>Sorbus aucuparia</i> (Rowan)	
222 Norway	$a + b \cdot D^2 + c \cdot D^2 \cdot H + d \cdot H^2 \cdot D + e \cdot H^2$
<i>Thuja plicata</i> (Western redcedar)	
223 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
224 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 40)^e$
<i>Tilia cordata</i> (Tei)	
225 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$
<i>Tsuga heterophylla</i> (Hemlock)	
226 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
227 Norway	$a \cdot H^b \cdot D^c \cdot (H - 1.3)^d \cdot (D + 100)^e$
<i>Ulmus</i> spp. (Elm, Orme, Ulm)	
228 Belgium	$a + b \cdot D + c \cdot D^2 + d \cdot D^3 + e \cdot H + f \cdot D^2 \cdot H$
229 Netherlands	$D^a \cdot H^b \cdot \exp(c)$
230 Romania	$a \cdot 10^{(b \cdot \log(D) + c \cdot \log(D)^2 + d \cdot \log(H) + e \cdot \log(H)^2)}$

**App. C**

	a	b	c	Parameters d	e	f	g
215	0.00095853	2.040672356	-0.04354461	0.56366437	-	-	-
216	$8.839 \cdot 10^{-5}$	1.8905	0.0469	0.8059	-0.0045	-	-
217	-0.011724	0.0000765	0.75	-	-	-	-
218	-1.86827	0.21461	0.01283	0.0138	-0.06311	-	-
219	0.00011585	1.6688	0.1090	0.7781	0.0269	-	-
220	$4.281 \cdot 10^{-5}$	2.0766	-0.1296	0.6843	0.2745	-	-
221	$7.325 \cdot 10^{-5}$	1.5598	0.0302	0.8572	0.1791	-	-
222	-1.86827	0.21461	0.01283	0.0138	-0.06311	-	-
223	1.67887	1.11243	-2.64821	-	-	-	-
224	1.3057	3.9075	1.9832	-2.3337	-1.3024	-	-
225	0.00004124	1.9302	0.0209	0.129	-0.1903	-	-
226	1.76755	1.37219	-3.54922	-	-	-	-
227	0.4291	2.6153	1.9145	-1.2696	-0.6715	-	-
228	-0.034716	0.004268168	-0.00013227	$-1.7667 \cdot 10^{-6}$	0.00016516	$3.8311 \cdot 10^{-5}$	-
229	1.94295	1.29229	-4.20064	-	-	-	-
230	$3.992 \cdot 10^{-5}$	2.1569	-0.0933	1.0728	-0.0708	-	-

